

# Effects of physical activity upon the liver

Roy J. Shephard · Nathan Johnson

Received: 5 March 2014 / Accepted: 14 October 2014 / Published online: 4 November 2014  
© Springer-Verlag Berlin Heidelberg 2014

## Abstract

**Purpose** To review the responses of the liver to acute and chronic physical activity and to summarize relationships between physical activity and liver health.

**Methods** A systematic search of HealthStar/Ovid from 1975 through June of 2013, supplemented by articles from other sources.

**Results** 351 of 8,010 articles identified by HealthStar/Ovid were supplemented by 92 other papers; after focussing, the review was reduced to 435 citations. Prolonged acute exercise reduces hepatic blood flow, stimulating hepatic glycogenolysis, gluconeogenesis and synthesis of some proteins; however, lipid metabolism shows little change. Glutathione depletion suggests oxidative stress. Enzymes affecting carbohydrate metabolism are up-regulated, and lipogenic enzymes are down-regulated. The main triggers are humoral, but hepatic afferent nerves, cytokines, reactive oxygen species, and changes in hepatic blood flow may all play some role. Regular aerobic exercise training improves

blood glucose control during exercise by increasing glycogen stores and up-regulating enzymes involved in gluconeogenesis and carbohydrate metabolism. Resistance to oxidant stress is generally increased by training. Lipogenic enzymes are down-regulated, and lipid metabolism is augmented. Modulations of insulin, insulin-like growth factor, glucagon and interleukin-6 may trigger the adaptive responses to training. Cross-sectional and longitudinal studies show that regular exercise can reduce hepatic fat, but the effect on circulating aminotransferases is unclear and the modality and dose of physical activity optimizing health benefits need clarification.

**Conclusions** Regular moderate physical activity enhances liver health. Adverse functional changes can develop if habitual activity is inadequate, and extremely prolonged competitive exercise may also be harmful, particularly under harsh environmental conditions.

**Keywords** Diabetes mellitus · Fatty liver · Hepatic blood flow · Hepatic clearance · Metabolic syndrome · Obesity · Oxidative stress · Steatosis · Steato-hepatitis · Ultra-marathons

Communicated by Nigel A.S. Taylor.

R. J. Shephard  
Faculty of Kinesiology and Physical Education, University of Toronto, Toronto, ON, Canada

R. J. Shephard (✉)  
PO Box 521, Brackendale, BC V0N 1H0, Canada  
e-mail: royjshep@shaw.ca

N. Johnson  
Faculty of Health Sciences, University of Sydney, Sydney, Australia

N. Johnson  
Boden Institute of Obesity, Nutrition, Exercise and Eating Disorders, University of Sydney, Sydney, Australia

## Abbreviations

ACC	Acetyl-coa carboxylase
ADP	Adenosine diphosphate
AKT	Protein kinase B
ALT	Alanine transaminase
AMP	Adenosine monophosphate
AMPK	Adenosine monophosphate kinase
ARFRP1	ADP-ribosylation factor-related protein 1
AST	Aspartate transaminase
ATP	Adenosine triphosphate
BCKDH	Branched-chain alpha-ketoacid dehydrogenase

cAMP	Cyclic adenosine monophosphate
CK	Cytokeratin
CoA	Coenzyme A
CT	Computerized tomography
DNA	Desoxyribonucleic acid
eIF2	Eukaryotic initiation factor 2
ERK	Extracellular signal-regulated kinase
FFA	Free fatty acid
GGT	Gamma glutamyl transferase
GLUT-2	Glucose-transporter-2
G protein	Guanine-nucleotide binding protein
GTP	Guanosine triphosphate
HAD	<i>B</i> -hydroxyacyl-dehydrogenase
HDL-c	High-density lipoprotein cholesterol
HISS	Hepatic insulin sensitizing substance
HMG-CoA	Hydroxymethylglutaryl-CoA
HSP	Heat shock protein
IGF-1	Insulin-like growth factor-1
IGFBP	Insulin-like growth factor binding protein
IL	Interleukin
IMTG	Intramycocellular triglyceride
JAK	Janus kinase
JNK	c-jun N-terminal kinase
LDH	Lactate dehydrogenase
MAPK	Mitogen-activated protein kinase
MDA	Malondialdehyde
mRNA	Messenger ribonucleic acid
NAFLD	Non-alcoholic fatty liver disease
NF- $\kappa$ B	Nuclear factor kappa-B
NO <sub>x</sub>	Mononitrogen oxides
PECPK	Phosphoenolpyruvate carboxykinase
PERK	Protein-kinase like endoplasmic reticular kinase
SCD-1	Stearoyl-CoA desaturase-1
sFasL	Soluble Fas ligand
SREBP-1c	Regulatory element-binding protein-1c
STAT	Signal transducer and activator of transcription
TRB3	Tribbles-related protein 3
VLDL	Very low density lipoprotein triglycerides
$\dot{V}O_{2max}$	Maximal oxygen intake

## Introduction

The liver is a major body organ that plays a central role in the regulation of carbohydrate and lipid stores, and ensures an adequate supply of metabolites for both vigorous physical activity and the synthesis of muscle and brain tissue (Wasserman and Cherrington 1991; Kjaer 1998; Wahren and Ekberg 2007; Fritsche et al. 2008). It also plays a vital role in metabolizing and/or excreting many unwanted

substances from the circulation. However, until recently there has been a relative dearth of literature describing the effects of physical activity upon the liver.

Interest in this question has arisen from the understanding that hepatic fatty infiltration (fatty liver) is independently associated with the metabolic syndrome, cardiovascular disease and type 2 diabetes mellitus, and from emerging evidence of an inverse association between physical activity and the risk of developing fatty liver. The latter topic has been the subject of other reviews, although with five exceptions (Eslami et al. 2009; Socha et al. 2009; Thoma et al. 2012; Keating et al. 2012; Musso et al. 2012), these have been unstructured. The contribution of physical inactivity to the development of fatty liver has been commonly acknowledged, and many authors have recommended exercise programmes or an increase of habitual activity as one element in therapy for this pathology (Table 1). Historically, this reflected a widespread view that lifestyle therapy, including regular exercise, could moderate fatty liver by assisting in weight loss, but more recently there has been a growing perception that regular physical activity in itself can exert beneficial effects upon the liver.

Much less is known about the homeostatic role of the liver during acute and chronic physical activity, even in healthy individuals. Available human studies have generally documented positive effects of acute and chronic exercise on liver glucose homeostasis and overall benefits of exercise training upon lipids and lipoprotein metabolism, including a reduction in hepatic fat infiltration (steatosis). These tissue-level observations have been amplified by rodent investigations showing an association between inadequate habitual physical activity and a down-regulation of the key hepatic enzymes associated with glucose and fat metabolism.

The present review examines gross and cellular responses of the liver to both acute and chronic physical activity, spanning the spectrum from physical inactivity to large volumes of vigorous exercise. The primary intention is to compile the collective research (from rodent to human, and tissue-level to molecular) concerning the acute and chronic effects of physical activity upon the healthy liver, in order to characterize the typical physiological responses of the liver to exercise. The review includes exercise-induced effects on hepatic blood flow and lipid and protein metabolism, with particular emphasis upon the interactions between exercise and glucose homeostasis, the molecular changes underlying these interactions, and their potential triggers. Secondary goals are to examine the patterns of regular physical activity associated with maintenance of normal hepatic function, and to document relationships between inadequate habitual physical activity and abnormalities of hepatic function.

**Table 1** Reviews considering the potential of exercise in the treatment of steatosis

Author	Number of references	Type of review	Comments	Conclusions
Alisi et al. (2010)	54	Unstructured review	Drugs might enhance response	Combination of diet and exercise can modify course of simple steatosis
Bacchi and Moghetti (2013)	18	Unstructured review		Exercise may reduce hepatic fat and hepatic enzymes, independent of dietary modifications
Caldwell and Lazo (2009)	48	Unstructured review		More exercise seems better, but optimal intensity not known
Cheung and Sanyal (2009)	101	Unstructured review	Some discussion of possible cellular mechanisms	Weight loss and increased exercise consistently associated with improved liver histology
Centis et al. (2010)	65	Unstructured review	Drugs should remain a second line of treatment	Weight loss and physical activity have specific therapeutic role
Conjeevaram and Tiniakos (2011)	26	Editorial review		Physical activity recommended, but lack of criteria for intensity, volume and duration of exercise
Della Corte et al. (2011)	96	Unstructured review	Drugs may serve as adjuvants	Weight loss diet and exercise are first line therapeutic measures
Dowman et al. (2011)	167	Unstructured review	No highly effective drug treatment	Need for multifaceted treatment: diet, exercise and behavioural counselling
Duvnjak et al. (2009)	151	Unstructured review	Value of pharmacological adjuvants unproven	Diet and exercise first line interventions, but problems of poor compliance. Lack of specific exercise guidelines
Eslami et al. (2009)	70	Systematic review		No effective treatment for steatosis- but weight loss and increased physical activity can improve enzymes and hepatic histology
Johnson and George (2010)	57	Unstructured review	Weight loss needs to be 3–10 % of body mass to reduce steatosis	Enzyme levels reduced and steatosis improved with exercise independent of weight loss
Keating et al. (2012)	47	Systematic review and meta-analysis to August 2011	12 studies included in meta-analysis	Beneficial effect of exercise on liver fat with little or no weight loss. No effect on ALT
Mencin and Lavine (2011)	95	Unstructured review	Vitamin E may be a helpful adjunct	Treatment remains diet and exercise
Musso et al. (2012)	140	Systematic review and meta-analysis	Statins and polyunsaturated fats also improved steatosis, but effects on liver histology unclear	Lifestyle weight loss >7 % safe, improved hepatic risk profile, but achieved by <50 % of patients
Nobili et al. (2011)	40	Unstructured review		Improved physical activity and nutrition needed to manage and treat condition
Nobili and Sanyal (2012)	72	Unstructured review	No drugs yet of proven value	Best approach is diet + exercise, but motivation is difficult
Rector and Thyfault (2011)	93	Unstructured review		Due to physical inactivity, low aerobic fitness and overnutrition singly or in combination
Reynoso and Lavine (2012)	10	Editorial	Conclusions based on guidelines of American Association for the Study of Liver Diseases, the American Gastroenterological Association, and the American College of Gastroenterology	Exercise (unspecified) might reduce steatosis but effect on other aspects of histology unknown
Rodriguez et al. (2012)	25	Semi-structured review		Optimal exercise regimen in terms of effectiveness and compliance has yet to be determined: reduction in sedentary time might also help

Table 1 continued

Author	Number of references	Type of review	Comments	Conclusions
Socha et al. (2009)	57	Structured review	Main focus on patients unable or unwilling to change lifestyle	Data insufficient to determine value of alternatives to lifestyle change
Spassiani and Kuk (2008)	28	Unstructured review	Most studies to date have included multiple therapies	Independent effect of exercise upon liver fat content unclear
Targher et al. (2010)	125	Unstructured review	Pharmacotherapy a questionable option	Team of experts needed to correct “unhealthy” habits, including exercise specialist
Thoma et al. (2012)	49	Structured review		23 studies identified; seven had control groups, six randomized. Consistent reductions in liver fat and ALT, correlated most strongly with decrease in body mass. Usually reduced insulin resistance, but changes of fibrosis less consistent
Yoneda et al. (2006)	Not stated in abstract	6-page review in Japanese		Since central obesity and insulin resistance are likely causes, exercise and nutritional counselling warranted. Lack of specific exercise guidelines
Zelber-Sagi et al. (2011)	151	Unstructured review	2-year compliance with physical activity as low as 20 %	Lifestyle modification with increased physical activity first line, aim of 5–10 % weight reduction

## Search techniques and classification of physical activity

The data base of HealthStar/Ovid was scanned from 1975 through June of 2013, using the terms exercise, exercise therapy, exercise training, activity, motor activity and physical activity vs. liver, liver cirrhosis, liver disease, fatty liver, liver failure, liver neoplasms, liver regeneration and liver transplants. This search yielded 8,010 hits. All articles that included a full abstract were examined, and 351 items that were specifically designed to examine the effects of acute or chronic physical activity upon the liver were included. This initial database was supplemented with 92 articles gleaned from reference lists and the authors' own extensive personal files; 8 reports were eliminated with a subsequent focusing of the review to exclude studies of the gall bladder and biliary tract, thus yielding a total of 435 citations.

For the purpose of this review, we have adopted the international consensus definitions (Bouchard and Shephard 1994) of physical activity (any bodily movement produced by the skeletal muscles and resulting in significant energy expenditure) and exercise (physical activity undertaken purposefully, with the intent of developing physical or physiological condition (e.g. cycling, treadmill running or athletic competition in humans and wheel/treadmill running in rodents)). Semantic descriptions have arbitrarily classed the intensity of effort as: low (30–50 % of  $\dot{V}O_{2max}$ ), moderate (50–65 % of  $\dot{V}O_{2max}$ ), vigorous or strenuous (>65 % of  $\dot{V}O_{2max}$ ), and exhausting (exercise pursued to exhaustion), based on the ranges previously described (Thompson 2010). Prolonged exercise is arbitrarily defined as bouts of 30 min or longer.

## Acute effects of moderate endurance exercise

### Hepatic blood flow

An acute bout of exercise transiently reduces blood flow to the liver. Estimates of hepatic blood flow in humans are commonly based upon indocyanine clearance. Studies using this technique have shown that blood flow to the liver and viscera decreases by up to 20 % during a brief period of vigorous effort, and even more if exercise is prolonged or is undertaken in a hot environment (Lundbergh and Strandell 1974; Rowell et al. 1964; Rowell 1974; Rowell 1986; van Wijck et al. 2011). There appears to be a dose–response relationship, with hepatic blood flow decreasing progressively as exercise intensity increases towards  $\dot{V}O_{2max}$ .

Elimination of indocyanine depends upon both hepatic blood flow and cellular function (Daemen et al. 1989). Thus, it has been argued that this method of measuring hepatic blood flow may be confounded by altered hepatic

metabolism during heavy exercise. Nevertheless, the trends indicated by the indocyanine method have been corroborated by studies based on sorbitol clearance; the latter technique has shown blood flow reductions of ~40 % at 40 % of  $\dot{V}O_{2\max}$  (Busse et al. 2004), of 60–70 % at 60–70 % of  $\dot{V}O_{2\max}$  (Kempe et al. 2000), and of 83 % during near-maximal exercise (Schoemaker et al. 1998). Although the magnitude of the immediate decrease in hepatic blood flow remains contentious (Froelich et al. 1988; Flamm et al. 1990), observations on human subjects have been confirmed by animal studies, where para-aminohippuric acid and sulfobromthalein were injected into a mesenteric vein, and samples drawn from both portal and hepatic veins (Katz and Bergman 1969; Yano et al. 1996).

On cessation of acute exercise, recovery of the resting hepatic blood flow appears to be rapid and indeed there is some evidence from ultrasound studies of human hepatic portal blood flow that hepatic blood flow exceeds normal resting levels for a few hours following physical activity (Hurren et al. 2011). This may reflect inflammation; arguably, it also serves to replenish glycogen reserves and speed the clearance of triacylglycerol from the circulation (Hurren et al. 2011).

We may thus conclude that although vigorous exercise induces a substantial immediate reduction of hepatic blood flow, this is rapidly reversed during recovery, and there is no evidence of subsequent harm to the liver.

### Carbohydrate metabolism

Human experimental studies using the stable isotope technique have demonstrated that liver glucose output is increased during exercise (Ahlborg et al. 1974). This serves to maintain blood glucose levels and contributes to the overall increase in the rate of glucose oxidation that is observed using indirect calorimetry. The rate of glucose oxidation is closely matched to work rate, (Bergstrom et al. 1967; Romijn et al. 1993), although even when exercising at ~50–85 % of  $\dot{V}O_{2\max}$ , liver-derived glucose contributes considerably less to the total energy requirement than the oxidation of skeletal muscle glycogen (Romijn et al. 1993).

The increased liver glucose output is partly a result of glycogenolysis, particularly during the first hour or more of sustained exercise (Kjaer 1998). However, the relative contribution of hepatic gluconeogenesis to total glucose output increases progressively as work duration is increased, and it accounts for some 50 % of glucose production during physical activity that is prolonged for more than one hour (Suh et al. 2007). Lactate (Shephard 1982; Wasserman and Cherrington 1991; Nielsen et al. 2007), amino acids (released from skeletal muscle through the action of cortisol), and glycerol all contribute to gluconeogenesis during exercise (Rowell 1971). As hepatic glycogen reserves become

depleted, the rate of gluconeogenesis is usually insufficient to sustain vigorous exercise, and a decline in the blood glucose concentration can therefore occur unless the work rate is reduced (Ahlborg et al. 1974). Depending upon an individual's training status and diet, both liver and muscle glycogen reserves can be almost completely exhausted over 90–180 min of vigorous aerobic exercise (Terjung et al. 1971).

In conclusion, exercise significantly increases liver glucose output by way of hepatic glycolysis and gluconeogenesis, making an important contribution to blood glucose control and oxidation during sustained endurance activity. These mechanisms can become exhausted during exercise such as a marathon run that continues for more than 90 min.

### Lipid metabolism

Under resting, fasted conditions, the liver accounts for a significant proportion (~40 %) of circulating fatty acid uptake, substantially exceeding the uptake of skeletal muscle (~15 %) (Jensen 1995; Meek et al. 1999). A portion of these fatty acids are converted to ketones or oxidized by the liver and other tissues in the splanchnic vascular bed (Havel et al. 1970; Wolfe et al. 1976). There is also a significant re-esterification of fatty acids to triglycerides in the liver (Klein et al. 1989); the triglycerides can then be secreted as very low density lipoprotein triglycerides (VLDLs).

The liver's dominant role in disposing of circulating (included ingested) fatty acids is suspended during exercise. Hormonal responses to exercise increase the net availability of circulating FFA during physical activity (Wolfe et al. 1990), but with the ensuing changes in blood flow distribution, the majority of these FFAs are directed to the contracting muscle; their oxidation accounts for most of the whole-body fat that is metabolized during exercise, although there is also a small contribution from intramyocellular triglyceride (IMTG)-derived fatty acids (Romijn et al. 1993). The splanchnic vascular bed accounts for less than 20 % of whole-body FFA uptake during exercise (Wolfe et al. 1990), but the relative partitioning of this uptake between hepatic oxidation and triglyceride synthesis remains unknown. Whilst it is believed that hepatic VLDL-derived fatty acids can be oxidized by skeletal muscle (Kiens 1993), the hepatic release of triglyceride in the form of VLDL remains unchanged during exercise (Børshiem et al. 1999). The current consensus is thus that VLDL makes a trivial contribution to whole-body fat metabolism during exercise (Helge et al. 2001).

Even a sustained bout of physical activity appears to have little immediate effect upon hepatic lipid metabolism. For instance, endurance-trained men showed no change in proton magnetic resonance spectroscopy estimates of hepatic triglycerides following 90 min of cycle ergometry at

65 % of  $\dot{V}O_{2peak}$  (Johnson et al. 2012). Likewise, a 60-min bout of cycle ergometry at 60 % of  $\dot{V}O_{2max}$  had no influence upon hepatic lipid metabolism in sedentary women (Magkos et al. 2009). One review also concluded that exercise had no effect upon the hepatic concentrations of total lipids, phospholipids and cholesterol in normally fed rats (Gorski et al. 1990). However, the high levels of circulating fatty acids induced by 60–90 min bouts of exercise led to an increase of hepatic triglycerides 3–4 h post-exercise in both mice (Hu et al. 2010) and human (Johnson et al. 2012) studies. A single 4-h bout of swimming also up-regulated hepatic stearyl CoA desaturase in rats and increased hepatic triglyceride content (Ochiai and Matsuo 2012). Further, exercise to exhaustion increased the hepatic content of the bound form of alpha-lipoic acid (lipoyl-lysine), an important co-factor for many mitochondrial proteins that are active in metabolism (Khanna et al. 1998). On the other hand, a single 3-h bout of exercise to exhaustion decreased the hepatic fatty acid synthase mRNA and enzyme activity induced by a high carbohydrate diet in both normal and diabetic (streptozotocin-treated) rats (Fiebig et al. 2001).

In conclusion, in contrast to its effects on glucose and protein metabolism, an acute bout of exercise has little immediate effect upon hepatic lipid metabolism and it may actually slightly increase hepatic triglyceride content. However, evidence (detailed later) showing up-regulation of hepatic enzymes and an overall reduction in hepatic fat levels with chronic exercise suggests that this is a transient response, with no detrimental effect upon the liver, and that a positive adaptation leading to a reduction of hepatic triglycerides occurs post-exercise and/or with chronic exercise.

### Protein metabolism

Sustained exercise can augment the hepatic synthesis of a number of proteins, including albumin and insulin-like growth factor binding protein (IGFBP). The latter binds IGF-1, allowing the growth hormone to act continuously upon the liver in paracrine fashion, producing more IGF-1.

Isotope infusion studies in humans have demonstrated increases in both the fractional (6 %) and the absolute synthesis (16 %) of albumin 6 h after completing a session of vigorous interval exercise (Yang et al. 1998). In rats, an increase in hepatic IGFBP-1 mRNA expression was also observed for up to 12 h following vigorous treadmill running; this response may serve to curtail an excessive muscle glucose uptake immediately post-exercise, thus preventing hypoglycemia (Anthony et al. 2001).

There is also evidence from the determination of arterial–hepatic venous differences in human subjects that the splanchnic uptake of alanine, synthesized and released by

the peripheral muscles, is increased 15–20 % during mild and moderate exercise (Felig and Wahren 1971). Presumably, this then serves for gluconeogenesis, a view that is supported by an increase of sweat nitrogen during exercise (Lemon and Nagle 1981).

In conclusion, a sustained acute bout of exercise can increase hepatic protein synthesis, but during prolonged activity, the liver also plays an important role in forming glucose from amino acids that are released from skeletal muscle.

### Triggers of changes in hepatic glucose metabolism during exercise

The classical view has been that changes in hepatic glucose metabolism with exercise are largely a consequence of the altered hormonal milieu. Thus, the exercise-induced increase in gluconeogenesis is stimulated by an attenuated secretion of insulin (Kjaer et al. 1993) and rising glucagon concentrations (Wasserman et al. 1989). If exercise is prolonged for more than one hour, these changes can be accentuated by declining plasma glucose concentrations (Trimmer et al. 2002) and depletion of hepatic glycogen reserves (Wahren et al. 1971; Peterson et al. 2004). Rising glucagon levels boost the extraction of glucose precursors from the blood, accelerating their conversion to glucose (Wasserman et al. 1989) and also stimulating glycogenolysis (Wasserman et al. 1995). The stimulation of glucagon receptors increases concentrations of cyclic adenosine monophosphate (cAMP), with activation of protein kinase A and extracellular signal-regulated kinase (ERK) (Jiang et al. 2001), and it also amplifies adenosine monophosphate kinase (AMPK) signaling (Berglund et al. 2009).

Somewhat surprisingly, moderate exercise does not cause much change in peripheral glucagon levels; however, this may be because plasma concentrations do not necessarily reflect glucagon levels in the portal vein (Wasserman et al. 1993). During vigorous exercise, catecholamine secretion may also play a regulatory role, either by providing the liver with additional substrate from adipose lipolysis and increased peripheral lactate formation (Wasserman et al. 1991), or by activating hepatic catecholamine receptors and thus mitogen-activated protein kinase (MAPK) (Christensen and Galbo 1983). Against this last hypothesis, hepatic glucose output does not seem to be affected greatly by adrenoceptor blockade (Coker et al. 1997).

Some correlate of glycogen depletion, albeit changes in concentration of a substrate, a derivative of substrate oxidation, an energy-related compound such as ATP, or an alteration in cell volume, might also trigger metabolic alterations more directly via the hepatic afferent nerves (Lavoie 2002). In support of this view, if glucagon secretion is suppressed,

an increased activity of the hepatic sympathetic nerves can be detected (van Dijk et al. 1994). On the other hand, glucose release is unaffected by hepatic nerve blockade (Kjaer et al. 1993; van Dijk et al. 1994), and hepatic denervation did not alter the glycemic response of rats to a brief bout of exercise (Lindfeldt et al. 1993).

Some of the changes seen during exercise may occur independently of both hormones and the autonomic nerves, with muscle-derived interleukin-6, for instance, playing a triggering role (Banzet et al. 2009). There are a number of pointers to an action of IL-6 upon the liver. For example, IL-6 stimulation of hepatoma cells increased their glucose production, and the injection of IL-6 into mice induced a small increase of hepatic phosphoenolpyruvate carboxykinase (PECPK) (Fritsche et al. 2010). Exercised mice also showed an increase of the hepatic chemokine CXCL-1 that attracts neutrophils and is involved in inflammation and wound healing, and muscle-derived IL-6 seems the trigger for this response (Pedersen et al. 2011). Finally, IL-6 may mediate the very large increase of hepcidin, a hormone that inhibits iron uptake, as seen in some athletes following prolonged and strenuous exercise such as a marathon run (Roecker et al. 2005; Peeling 2010).

Exercise might also modify liver function through an increased generation of reactive oxygen species, much as in skeletal muscle (Davies et al. 1982; Koyama et al. 1999; Liu et al. 2000). Certainly vigorous prolonged exercise (particularly if performed under hot and humid conditions) significantly restricts visceral blood flow (Wade and Bishop 1962; Rowell 1971), temporarily depriving the internal organs of an adequate oxygen supply (Shephard 2013), and this could favour the formation of reactive oxygen species. The exercise-induced up-regulation of heat shock proteins in rat studies seems to support this hypothesis (Salo et al. 1991; Kregel and Mosely 1996; Gonzalez and Manso 2004). On the other hand, some researchers have found little evidence of oxidative stress in the rat liver following acute exhausting exercise (Bejma et al. 2000; Ogonovszky et al. 2005), with no changes in the activity of anti-oxidant enzymes (Hoene and Weigert 2010).

There seems no fundamental reason why triggers should differ between humans and laboratory animals, but one issue to remember in interpreting these various findings is that much of the available research has been conducted on rodents, where hepatic glycogen reserves are relatively much larger than in humans (Baldwin et al. 1973; Terjung et al. 1974).

In conclusion, there remain several competing hypotheses concerning triggers to the hepatic adaptations of carbohydrate metabolism during acute exercise. It is unclear whether changes in hormonal milieu, substrate/metabolite concentration, cytokines, reactive oxygen species or associated changes in hepatic blood flow initiate these metabolic changes; further research is needed to decide among these possibilities.

## Molecular changes

Information on the molecular changes induced in the liver induced by acute exercise is based almost exclusively on studies of normally inactive rodents (Table 2). An analysis of the hepatocyte transcriptome in mice following 60 min of moderate intensity exercise showed that 352 transcripts were up-regulated, and 184 were down-regulated. Many of these changes affected genes that are important for glycolysis, gluconeogenesis and fatty acid metabolism (Hoene and Weigert 2010). Some of these same genes were activated in skeletal muscle, but the response was generally more marked in the liver. The effect was also transient, disappearing within a few hours of ceasing exercise (Hoene and Weigert 2010; Hoene et al. 2010).

Exercise has consistently led to an up-regulation of gluconeogenic and metabolic enzymes such as glucose-6-phosphatase, pyruvate dehydrogenase, and phosphoenolpyruvate carboxykinase (PEPCK) (Banzet et al. 2009; Hoene et al. 2009), reduced expression of lipogenic enzymes (Griffiths et al. 1996; Fiebig et al. 2001), the induction of metabolic regulators such as insulin receptor substrate (Hoene et al. 2009), and an up-regulation of the genes induced by energy depletion. Cortisol is normally implicated in the activation of hepatic PEPCK transcription. Such an involvement is supported by the greatly attenuated exercise responses in adrenalectomized animals, and by the absence of an exercise effect in transgenic mice with deletion of the glucocorticoid regulatory unit (Friedman 1994).

Another exercise-related change is activation of interleukin-6 type cytokine/cytokine receptor signaling, particularly the Janus kinase (JAK)/signal transducer and activator of transcription (STAT) pathway, which can transmit information from outside of the cell to gene promoters on intracellular DNA (Hoene and Weigert 2010). Exercise was shown to increase the activity of hepatic AMP-activated protein kinase activity several fold in rats and this response was diminished in IL-6 knock-out animals (Kelly et al. 2004); however, it is not yet quite clear how this particular cytokine is involved, since IL-6 deficiency did not impair the induction of metabolic genes by moderate exercise (Fritsche et al. 2010). An acute exercise bout also induced a marked activation of the mitogen-activated protein kinase (MAPK) signaling pathway, which transmits information from receptors on the cell surface to nuclear DNA (Hoene et al. 2010).

There is an up-regulation of the hepatic p53 tumour-suppressing pathway that guards against genome mutation (Hoene and Weigert 2010; Hoene et al. 2010), much as seen in skeletal muscle, particularly following eccentric exercise (Chen et al. 2002; Hoene and Weigert 2010). Exercise also stimulates an increased synthesis of heat shock

**Table 2** Molecular changes in the liver induced by acute exercise

Author	Change	Biological significance
Anthony et al. (2001)	Increased IGFBP-1 mRNA expression	May help to limit hypoglycaemia post-exercise
Dohm et al. (1985); Nizielski et al. (1996); Banzet et al. (2009); Hoene et al. (2009)	Up-regulation of glucose-6-phosphatase, pyruvate dehydrogenase, phosphoenol pyruvate carboxylase	Increase of gluconeogenesis
Griffiths et al. (1996); Fiebig et al. (2001)	Reduced expression of lipogenic enzymes	Less fat synthesis
Hoene et al. (2009)	Induction of insulin-receptor substrate	Increases insulin binding, facilitates gluconeogenesis
Hoene and Weigert (2010)	Changes in 352 gene transcripts	Many of genes active in glycolysis, gluconeogenesis and fatty acid metabolism
Friedman (1994); Ropelle et al. (2009)	Increase of PECPK gene and PECPK mRNA in mice, decrease of PECPK and glucose phosphatase in obese rats	Regulation of gluconeogenesis
Huang et al. (2007)	Increase of adiponectin receptor 1, decrease of adiponectin receptor 2, increase of the transcription factor forkhead box O1	Regulation of gluconeogenesis
Heinrich et al. (2003); Banzet et al. (2009); Hoene et al. (2009)	Increase of peroxisome proliferator-activated receptor coactivator PGC-1	Increased levels of protein that regulates mitochondrial biogenesis
Gonzalez and Manso (2004)	Increased synthesis of heat shock protein molecular chaperones	Protection of proteins against environmental stressors
Hoene and Weigert (2010); Hoene et al. (2010)	Activation of the mitogen-activated protein kinase (MAPK) signaling pathway and p53 tumour suppressing pathway	Regulation of proliferation, gene expression, differentiation and cell survival
Kelly et al. (2004)	Increased hepatic AMP-activated protein kinase activity	Regulation of carbohydrate and lipid metabolism; effect less in IL-6 knock-out mice
Haase et al. (2011)	Increased mRNA and cytochrome <i>c</i> protein	Improves oxidative capacity
Hansen et al. (2011)	Increased production of follistatin	Inhibits myostatin, facilitates muscle hypertrophy
Khanna et al. (1998)	Increased hepatic content of bound form of alpha-lipoic acid	Co-factor for many mitochondrial proteins active in metabolism
Lavoie et al. (2002); Leu and George (2007)	Increased blood levels of insulin binding growth factor binding protein	Helps glucose regulation by neutralizing insulin-like effects of insulin-like growth factor-1; also counters apoptotic effect of p53
Ochiai and Matsuo (2012)	Up-regulation of hepatic stearyl CoA desaturase	May protect against insulin resistance
Roecker et al. (2005); Peeling (2010)	Increase of hepcidin	Inhibition of iron uptake; could contribute to athlete's anaemia



protein (HSP-72, HSP-73 and of the glucose-regulated proteins 75 and 78) in rat liver (Gonzalez and Manso 2004) with a marked up-regulation of genes associated with a stress response, such as transcription factors of the Fos/Jun-family (Hoene et al. 2010). The only human study available to date confirmed that HSP72 from the hepato-splanchnic viscera increased in the first 2 h following prolonged moderate intensity exercise (Febbraio et al. 2002).

Finally, increased production of the transforming growth factor follistatin has been demonstrated in mouse liver during exercise (Hansen et al. 2011). Follistatin inhibits myostatin, thus encouraging muscle hypertrophy, and it also acts as a growth promoter for hepatocytes (Fuwii et al. 2005). Observations on human subjects during cycling have confirmed that liver is the source of the follistatin during exercise; active muscles do not liberate this substance into the circulation (Hansen et al. 2011).

In conclusion, information on the molecular changes during acute exercise generally conforms with expectations based on the gross biochemical changes, including the up-regulation of metabolic enzymes and a decreased expression of lipogenic enzymes. There is also an up-regulation of systems protecting against gene mutation and heat shock, and an increased formation of transforming growth factors such as follistatin.

### Adverse responses to acute exercise

Whilst moderate exercise appears to have little effect upon either the morphological characteristics of hepatic tissue (Latour et al. 1999) or liver function and oxidative stress, histological changes, impaired pharmacokinetics, markers of oxidative stress and altered blood levels of hepatic enzymes have been observed after heavy and prolonged exercise (Table 3, particularly if there is associated heat stress (Berg and Keul 1982; Hassanein et al. 1992; Giercksky et al. 1999; Eran et al. 2004; Miura et al. 2010). Such findings all point to adverse changes of hepatic function, which could have negative implications for those participating in prolonged endurance events such as marathon running, distance cycling, and long-distance triathlons. However, information to date suggests that normal liver function is regained, at most within a few days of ceasing exercise.

### Histological changes

The histological changes associated with vigorous and/or exhausting exercise have been examined mainly in animals. A very heavy bout of exercise has been shown to cause an inflammatory response, with an increase in peripheral leucocyte count (Kayashima et al. 1995). Exercise that results in hepatic hypoxia can also predispose to central lobular

necrosis (Rowell 1971; Prapatsorn et al. 2010). Prolonged exercise to exhaustion has further been shown to induce mitochondrial swelling in hepatocytes surrounding hepatic venules, and oncotic and/or apoptotic necrosis of the hepatocytes in rodents (Yano et al. 1997; Huang et al. 2013).

### Impaired pharmacokinetics

The study of pharmacokinetic changes induced by exercise provides further insight into possible disturbances of liver function and health. Most pharmacokinetic investigations have been conducted in human subjects. The elimination of “low clearance” drugs such as acetaminophen, antipyrine, diazepam, amylobarbitone and verapamil is affected primarily by changes in hepatic enzyme activity and biliary excretion (Khazaenia et al. 2000). Their clearance is largely unaffected by either moderate exercise (Balasubramian et al. 1970; Swartz et al. 1974; Klotz and Lucke 1978; Mooy et al. 1986; Loniewski et al. 2001) or very prolonged low to moderate intensity activity such as 6–9 h of marching (Theilade et al. 1979; Fabbri et al. 1991). In contrast, bouts of vigorous and/or prolonged exercise reduce the elimination of such substances as indocyanine, bromsulphthalein, sorbitol and lidocaine (Mooy et al. 1986; van Griensven et al. 1995), whose clearance rate mainly reflects hepatic blood flow (Rowell et al. 1964; Døssing 1985). There remains a need to examine the effects of vigorous and very prolonged events such as ultra-marathon runs or long course triathlon on hepatic clearance function, particular with respect to low clearance drugs.

### Oxidant stress

As with the gross changes observed in lipid and lipoprotein concentrations, significant changes in oxidant status may develop after, rather than during an exercise bout (Koyama et al. 1999). It is therefore important that research studies continue observations sufficiently far into the recovery period to detect such changes. Importantly, as in skeletal muscle, these changes are not necessarily ‘adverse’ per se, as they may be important factors in inducing adaptations to regular exercise (Hoene and Weigert 2010).

Some (Neubauer et al. 2008; Pinho et al. 2010; Turner et al. 2011) but not all studies (Margaritis et al. 1997) in humans have demonstrated a transient increase of oxidant stress following prolonged and/or vigorous exercise. However, none of these studies have examined changes within the liver itself.

Animal studies (Table 4) provide more direct evidence that exhausting exercise causes oxidative stress in the liver. For instance, the hepatic glutathione levels have been shown to fall in rats following exhausting exercise, reflecting a large increase in oxidative metabolism, and a reduced ability to buffer reactive oxygen species (Sen

**Table 3** Reports of adverse changes in the liver following prolonged endurance exercise

Author	Subjects	Exercise	Findings	Comment
Apple and McGue (1983)	Two male runners	6 weeks of training for a marathon	ALT increased in 1/2 subjects; large increases of LDH and CK	
Beard et al. (1979)	Two joggers with heat stroke		Reduced level of clotting factors produced by liver	
Bunch (1980)	Six runners		Clinically “abnormal” levels of hepatic enzymes	
Böger-Mendonça et al. (2008)	Six male athletes	Half-triathlon	Comparison of blood samples before and after race; significant change in AST and ALP but not ALT	Moderate temperatures. All enzyme values remained within normal limits
De Paz et al. (1995)	13 male runners, mean age 36 years	100-km race	Post-race increases in ALT +42 %, AST +193 %, GGT +56 %, CK +2,000 %	Cool conditions; serum bilirubin +106 %. Also decrease of serum haptoglobins
Fallon et al. (1999)	Seven male, two female runners	1,600-km ultramarathon	Increases of ALT, AST, GGT, LDFH, CK; ALT remains high when AST and CK falling	Temperatures 11–32 °C
Foigt et al. (1976)	Six male volunteers	Cycle ergometer exercise at 70–85 % maximal oxygen intake to exhaustion (26–60 min)	Hepatic vein shows increased content of ALT	
Hammouda et al. (2012)	18 male football players	Wingate test	Small increases of CK 11 %; AST 10 %; ALT 16 %; LDH 13 %	
Holly et al. (1986)	Six male, three female triathlon competitors	Hawaiian Ironman competition	SGOT +700 %, SGPT +262 %, LDH +222 % immediately after competition	Initial resting values high normal; enzymes marginally increased 5–6 days later
Kayashima et al. (1995)	14 soldiers aged 24–36 years	80-km trek on limited diet over 4 days	Enzymes sampled immediately and after 8 days. Increases: AST 177, 152 %; ALT 39, 234 %; LDH 66, 8 %; CK 208, –68 %	Associated leucocytosis and bilirubinaemia immediately after exercise
Kratz et al. (2002)	32 male, five female runners, average age 49 years	Boston marathon	AST +265 %, ALT +37 %, CK +2,343 % increased after race, 24 h > 4 h	Cool environment; bilirubin increased 60 %, serum urea nitrogen 29 %
Lippi et al. (2011)	15 healthy males	21-km half-marathon	AST, LDH, CK increased 0–24 h following run	ALT not measured
Litvinova and Viru (1995)	Male Wistar rats aged 10–12 weeks	10-h swimming with loading 10 % body mass	103 % increase in hepatic <sup>14</sup> C urea content	Effect decreased by adrenalectomy
Long et al. (1990)	Ten athletes	Short triathlon (10-km run, 20–40 km cycle, 1.0–1.5 km swim)	Modest increases of AST (30 %) and LDH (53 %)	Moderate temperatures; ALT not measured
Malinoski (1992)	Three soldiers	Several days of strenuous training	Increases of ALT, AST and CK	
Mena et al. (1996)	Professional cyclists	800 km/6 days and 2,700 km/20 days with overnight rests	Increases of ALT, AST, ALP, and (in longer race) LDH; partial return to normal with overnight rest	Cumulative increase over race
Nagel et al. (1990)	55 runners	1,000 km in 20 days	AST increased 500 %, ALT 300 %, GGT 600 %, CK 2,000 %; ALT remains high when AST and CK falling	Decreases in serum albumin and choline esterase (could reflect decreased synthesis or increase of IL-1)
Nathwani et al. (2005)	Four prison inmates	Three, vigorous squatting; one, long-distance run	Increases of AST, ALT, CPK and LDH	Liver damage unlikely since normal serum bilirubin and prothrombin times

Table 3 continued

Author	Subjects	Exercise	Findings	Comment
Noakes and Carter (1976)	13 athletes	160-km run	Increases of LDH 241 %; AST 821 %; CPK 1,732 % in those completing event	Temperatures not stated; total bilirubin also increased twofold; ALT and GGT not determined
Noakes and Carter (1982)	18 experienced, five novice competitors	56-km ultramarathon	Greater rise of AST and CPK in novices, despite slower running speed	
Ohno et al. (1988)	Seven sedentary male students	Running 5 km, 6 times/week for 10 weeks	50 % decrease of resting GGT	
Pettersson et al. (2008)	15 healthy men not used to weight-lifting	1 h of weight-lifting	AST, ALT, LDH, CK all remained elevated for 7 days post-exercise	Laboratory conditions
Rama et al. (1994)	7 well-trained male distance runners, mean age 37 years	100-km road race	GGT + 19 %, CK +3,121 %	Cool conditions; ALT not measured
Richard's et al. (1979)	43 successful runners (28 M, 16 F) vs. ten who collapsed (9 M, 1 F)	14 km Sydney city to surf run	Casualties showed higher values for blood urea nitrogen, serum creatinine, uric acid and bilirubin	
Shapiro et al. (1973)	26 untrained men	110 km march in 2 days	Increases in CK, GOT and aldolase	Midday temperatures 30 °C; ALT not measured. Increased enzyme levels only seen in those marching at 6 km/h
Smith et al. (2004)	27 male, seven female runners, aged 18–65 years	Marathon run	Significant increases in CK, AST, LDH immediately following event	ALT not determined; climate not specified
Suzuki et al. (2006)	Nine well-trained male triathletes	Ironman triathlon	Significant increases immediately and especially 24 h after race: ALT 185 %; AST 759 %; CK 2,680 %; GGT –20 %	Moderately warm conditions
Takahashi et al. (2007)	Seven male rugby players, average age 21 years	Two successive Rugby sevens matches of 10 min duration, with 4-h interval	Increases of CK 42 %; LDH 25 %; AST 13 %; but not ALT	Cool conditions
Van Rensburg et al. (1986)	23 male athletes, average age 33 years	Triathlon competition	Significant increases of AST, CK and LDH immediately post-race	ALT not measured; 4.5 % decrease of body mass over event
Waskiewicz et al. (2012)	14 male runners mean age 43 years	24-h ultra-marathon	Increased enzymes 24 h after run; ALT 350 %, AST 1,354 %, CK 1,204 %, no change of GGT	Cool conditions
Wu et al. (2004)	Ten males, one female	24-h marathon	Blood enzymes tested immediately, 2 and 9 days after event. AST 1,344, 630, –8 %; ALT 237, 259, 44 %; LDH 286, 205, 59 %; no significant change of GGT	Moderately warm conditions; bilirubin increased immediately post-race

et al. 1992). Increased blood levels of malondialdehyde (MDA, a marker of lipid peroxidation), NO<sub>x</sub>, and xanthine oxidase have also been shown in mice, accompanied by large increases in serum aminotransferases and lactate dehydrogenase (LDH) (Kayatekin et al. 2002). Relative to control animals, liver samples showed increased neutrophil infiltration and reductions in levels of superoxide dismutase, catalase and glutathione peroxidase (Huang et al. 2008). Other studies of rats have found significant increases in measures of hepatic lipid peroxidation following exhaustive exercise (Turgut et al. 2003; Aydin et al. 2005). There is also a substantial rise in the temperature of hepatic tissue during exhausting exercise, and rodent studies have demonstrated an associated increase in concentrations of 70 and 72 kDa HSPs (Salo et al. 1991; Gonzalez and Manso 2004).

Thus, animal studies support the human inference of increased oxidant stress and increased hepatic concentrations of HSP following heavy exercise, but further research is needed to determine how far such changes are an inherent part of the adaptive response to physical activity.

#### Serum hepatic enzyme levels

Clinicians frequently evaluate human hepatic function in terms of serum levels of hepatic enzymes. Short periods of acute exercise usually have little or no effect upon such indices (Takahashi et al. 2007; Hammouda et al. 2012). However, in the hours following vigorous and very prolonged exercise such as marathon or triathlon competition, many investigators have found increased concentrations of serum aminotransferases, often accompanied by increased bilirubin concentrations and markers of inflammation such as IL-6 and C-reactive protein, similar to the findings in laboratory animals after prolonged vigorous exercise (Moses 1990; Prapatsorn et al. 2010) (Table 3). The cause of these changes (recent physical activity, hepatic injury, haemolysis, or muscle injury) remains unclear (Kindermann et al. 1983; Koutedakis et al. 1993; Rosales et al. 2008). Confirmation of sustained hepatic malfunction has been sought in decreased plasma levels of albumin, globulin and cholinesterase (Nagel et al. 1990; Wu et al. 2004), although reduced concentrations of these substances could also reflect the influence of increased serum concentrations of interleukin-1 (Nagel et al. 1990). Further information is thus required before we can interpret exercise-induced changes in clinical liver function tests as evidence of hepatic damage.

#### Summary of responses to acute exercise

Hepatic responses to acute exercise include a decrease in regional blood flow, and an increase of glucose output

by way of glycogenolysis and gluconeogenesis. These changes are exacerbated as the intensity and duration of exercise is increased, and they contribute to maintenance of a stable blood glucose concentration. At rest, the liver is a major site for fatty acid uptake, much of which is re-packaged and secreted as VLDLs. However, during an acute bout of physical activity, possibly as a consequence of reduced hepatic blood flow and increased fatty acid uptake by muscle, the liver adopts a more 'passive' role, with no measurable change in liver triglyceride concentrations. Albumin and IGF-1 levels are increased after an acute bout of exercise; they likely have growth-promoting effects and contribute to euglycaemia. Although glucagon and insulin have some regulatory influence, the precise stimuli triggering the changes in hepatic function that maintain blood glucose during exercise remain unclear, as do the underlying molecular adaptations. Changes in catecholamines, intracellular high-energy phosphate concentrations related to substrate availability, reactive oxygen species, cytokines and tissue hypoxia have all been suggested as playing a regulatory role. Understanding these stimuli and the molecular effects of acute exercise is made difficult by a lack of direct human evidence, and potential difficulties in translating rodent findings to an interpretation of human responses. Whilst moderate exercise appears well tolerated by the liver, vigorous and/or prolonged or exhaustive activity may result in inflammation, altered pharmacokinetics, oxidative stress and increases in concentrations of HSPs and serum amino-transferases. Vigorous and/or prolonged exercise can cause a slowing in the elimination of markers dependent upon hepatic blood flow, signs of oxidative stress in both humans and animals, and a transient appearance of hepatic enzymes in the serum. However, there is little evidence of permanent hepatic damage; such disturbances seem transient and possibly contribute to exercise adaptations.

#### Chronic effects of moderate endurance exercise

As with the acute effects of exercise, we will consider changes in the metabolism of carbohydrates, lipids and protein, and triggers for these changes. We will focus particularly upon the role of oxidant stress, and implications for hepatic function.

#### Carbohydrate metabolism

It is well known that regular exercise training increases a person's ability to sustain a higher work-rate during prolonged activity, and to exercise for longer before the onset of fatigue. One component of this change is an enhanced resistance to hypoglycemia during exercise. This is partly a

consequence of an increased capacity for skeletal muscle to store glycogen and to oxidize fat at the expense of glucose. Although there is relatively little human data, further ‘glucose sparing’ adaptations likely include an increased resting liver glycogen concentration and a reduced rate of both glycogenolysis and gluconeogenesis at any given intensity of exercise (Coggan et al. 1995; Murakami et al. 1997). Other changes associated with exercise training include a reduced availability of gluconeogenic precursors (lactate and glycerol) at a given volume of exercise, and altered hormonal responses (a higher insulin, and lower glucagon and catecholamine concentrations).

Rodent investigations generally agree with human observations in showing that the glyconeogenic and the gluconeogenic responses to glucagon are enhanced with training (Podolin et al. 2001; Drouin et al. 2004). However, there are some differences in the responses of rats, probably related to the fact that gluconeogenesis accounts for some 20 % of glucose production when humans undertake moderate exercise, whereas in rats the figure ranges from 40 to 70 %. In particular, training increases exercise hepatic glucose clearance in humans, but not in rats (Coggan et al. 1995).

Underlying mechanisms apparently include a normalizing of the ratio of inhibitory to stimulatory guanine-nucleotide binding protein (G protein), and a resultant increase in activity of the “second messenger” adenylyl cyclase (Podolin et al. 2001). The increased capacity for glucose output contributes to the ability of trained individuals to sustain higher work-rates and to maintain euglycaemia during exercise (Donovan and Sumida 1990). Moreover, the liver of a trained person has an increased absolute capacity for lactate (Donovan and Pagliassotti 1990) and alanine (Sumida and Donovan 1995) clearance, and associated gluconeogenesis (Sumida and Donovan 1993).

Most, but not all (James and Kraegen 1984), rodent studies have shown increases in activity of the enzymes and signaling molecules involved in both carbohydrate and lipid metabolism following aerobic training (Colombo et al. 2005; Aoi et al. 2011).

In conclusion, aerobic training induces metabolic adaptations in both humans and laboratory animals that help to conserve glucose homeostasis during prolonged exercise, including greater glycogen storage in both liver and muscle, and the sparing of carbohydrate through greater fat metabolism.

### Lipid metabolism

The enhanced ability to utilize fat during exercise following regular training is largely a function of adaptations in skeletal muscle (and associated hormonal changes); there is little evidence that the liver contributes to this response. This is perhaps understandable, given the apparently trivial role

of the liver in contributing to fat oxidation (via VLDLs) (Helge et al. 2001). Moreover, exercise training blunts the lipolytic hormone response to exercise, so that after training circulating concentrations of insulin and insulin-like growth factor binding protein-1 tend to be higher (Prior et al. 2012), and blood glycerol and FFA concentrations are lower at a given absolute exercise intensity (Martin et al. 1993). These changes further reduce the liver’s role, including its exposure to FFAs. Nevertheless, regular exercise is associated with alterations in lipid/lipoprotein metabolism, and it appears to reduce the amount of triglyceride stored in the liver.

Several studies have examined the effect of regular exercise upon liver fat content (discussed later), and the associated liver mass in rats and mice (Table 8). Most investigators have observed a reduction in liver mass with regular exercise training, although such findings have not been universal (James and Kraegen 1984; Murakami et al. 1997). A limitation of many animal studies is that control animals have lived unnatural lives of physical inactivity and over-eating relative to their natural state, and in consequence differences in hepatic tissue mass between sedentary and exercised animals have varied between investigations. For example, in the study of Yiamouyiannis et al. (1992), rats that were fed ad libitum and given free access to a running wheel also ate more, thus presenting with increased values for total, mitochondrial and cytosolic protein (Yiamouyiannis et al. 1992). The total activity of several enzymes was also increased, although the activity per g of liver or per g of hepatic protein remained unchanged (Yiamouyiannis et al. 1992).

The cardio-protective benefit of regular exercise in modifying circulating lipids and lipoproteins is well documented in both human subjects and experimental animals. Cross-sectional research shows that high-density lipoprotein cholesterol (HDL-c) levels are higher in regular exercisers versus their inactive counterparts (Williams et al. 1981), and HDL-c increases with exercise training interventions (Kelley et al. 2005, 2006; Dressendorfer et al. 1982; Terao et al. 1989). Similarly, exercise training may reduce circulating triglycerides and VLDL secretion (Tsekouras et al. 2008). These benefits are associated with a decreased activity of hepatic lipase (Thompson et al. 1991) and alterations in the levels of other hepatic enzymes involved in HDL-c remodelling (including cholesteryl ester transfer protein and lecithin cholesteryl acyl transferase) (Kraus et al. 2002; Halverstadt et al. 2007).

Inter-individual human differences in lipid responses to training programmes have been traced to a polymorphism in the hepatic lipase gene LIPC -514C-T (Brinkley et al. 2011). However, the relative contribution of acute versus chronic responses to these exercise-induced improvements in lipids and lipoproteins remains unclear (Cullinane et al. 1982; Thompson et al. 2001; Magkos et al. 2007).

**Table 4** Effects of chronic exercise upon oxidative stress in the liver

Author	Subjects	Training	Outcome
Burneiko et al. (2006)	60-day-old male Wistar rats	Swimming up to 60 min/day, 2 times/week for 8 weeks	Hepatic superoxide dismutase increased, catalase decreased, increase of total antioxidant substances
Chapados and Lavoie (2010)	6-week-old female Sprague–Dawley rats	Treadmill running for 6 weeks (progressing to 60 min/session at 26 m/min, 10 % slope), 5 times/week	Increased gene expression of unfolded protein response markers may protect against endoplasmic reticulum stress
da Silva et al. (2009)	3-month-old male mice	Intermittent or continuous or downhill running 45 min/day for 8 weeks increasing to speed of 16 m/min	Continuous or downhill running increased hepatic superoxide dismutase, decreased catalase, decreased lipoperoxidation and protein carbonylation
Gore et al. (1998)	4-month female Sprague–Dawley rats	10-week heavy training (2 h/day running 25 m/min up 10 % grade)	58 % increase of hepatic superoxide dismutase
Güündüz et al. (2004)	9-month-old albino Wistar rats	1 year of swimming 1 h/day	Reversed age-related decrease of hepatic catalase
Haase et al. (2011)	Mice	5 weeks of exercise (combination of 1 h/day of treadmill running, 14 m/min 10 % slope, and wheel running 5 days/week)	No change of mRNA expression for hepatic anti-oxidant enzymes
Ho et al. (2001)	2-month-old female Sprague–Dawley rats	3 months swimming (2 h/day)	Down-regulation of cytosolic aconitase, possibly counters increase production of NO and a related decrease in the iron content of the liver
Hong and Johnson (1995)	Normotensive and hypertensive male rats	10-week treadmill training, progressing to 60 min/day at 20 m/min, 5 % slope (exercise caused some weight loss)	Down-regulation of hepatic superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase, BUT decrease of TBARS seen 7 days post-training
Huang et al. (2010)	6-week-old Sprague–Dawley rats	12-week running up 12 % grade, 60 min/day	Increased anti-oxidant content of liver
Kiraly et al. (2010)	6-week-old male ZDF diabetic fatty rats	10-week voluntary wheel running	Decreased markers of inflammation and oxidative stress in the liver, lower circulating IL-6 and malondialdehyde levels, less hepatic protein oxidation, decreased c-Jun-terminal kinase activity (heralds apoptosis), lower PECPK levels, lower Ser (307)-phosphorylated insulin receptor substrate-1 concentrations
Navarro et al. (2003)	28-week-old mice	50-week treadmill running (5 min at 6–12 m/min, thrice weekly)	Reduced oxidative stress markers in hepatic mitochondria
Radak et al. (2004)	30-month-old rats	8-week exercise (2 h/day at 2.1 km/h)	Reversed age-related increased concentration of reactive oxygen species in the liver; attenuation of DNA binding activity of NF- $\kappa$ B (could decrease transcription of inflammation-related genes)
Reddy Avula and Ferandes (1999)	3-month-old mice	8-week treadmill running (45–50 min/day at 1 km/h)	Increased levels of hepatic catalase and glutathione transferase activity and lower lipid peroxide levels
Romani et al. (2009)	8-week-old male Sprague–Dawley albino rats	10-week treadmill running (1 h/day, 3 times/week at speeds increasing to 65 % of maximal oxygen intake)	Increased paraoxanase-3 (increases anti-oxidant properties of HDL-cholesterol)
Sen et al. (1992)	10-week-old Han Wistar rats	8-week treadmill training (2 h/day at 2.1 km/h)	21 % increase of hepatic glutathione levels
Sen et al. (1992)	15-week-old female beagle dogs	55 weeks of treadmill training (40 km/day at 5.5–6.8 km/h, 15 % slope)	30 % increase in activity of hepatic glutathione transferase S activity, 8 % increase of total hepatic glutathione

Table 4 continued

Author	Subjects	Training	Outcome
Venditti and di Meo (1997)	12-month-old rats	10-week swimming	Reduced lipid peroxidation and increased glutathione peroxidase, glutathione reductase and overall anti-oxidant activity
Wilson and Johnson (2000)	11–12 week male Sprague–Dawley rats	10-week treadmill running (60 min/day, 20 m/min 5 % slope)	Hepatic superoxide dismutase mRNA down-regulated, but catalase mRNA up-regulated

Rodent investigations have provided insights into the molecular changes underlying the effects of regular physical activity upon lipid/lipoprotein metabolism. Training sessions reduced hepatic acetyl-coenzyme A carboxylase and fatty acid synthase activity and mRNA (Askew et al. 1975; Fiebig et al. 1998, 2001, 2002; Lavoie and Gauthier 2006). Regular exercise also down-regulated the hepatic gene and protein content of stearoyl-CoA desaturase-1 (SCD-1), the rate limiting enzyme in the biosynthesis of saturated-derived monounsaturated fats that are a major constituent of VLDLs. Further, there was a down-regulation of the microsomal triglyceride transfer protein that plays a key role in the assembly and secretion of VLDL lipoprotein (Chapados et al. 2009), and training increased levels of hepatic mRNA for the ATP-binding cassette transporter A-1 that plays a vital role in membrane transport and plasma HDL cholesterol remodeling (Ghanbari-Niaki et al. 2007). Changes in the composition of hepatic phospholipids following training likely have implications for membrane properties, cell signalling and gene expression (Petridou et al. 2005).

In conclusion, exercise training increases muscular oxidation of fat and leads to molecular changes of hepatic function that reduce liver fat content and enhance the blood lipid profile.

#### Protein metabolism

An expansion of plasma volume is a well-recognized adaptation to regular exercise; expression of the hepatic albumin gene mRNA facilitates this response by increasing serum albumin concentrations, and rodent studies indicate that such an adaptation can occur within days of the initiation of training (Bexfield et al. 2009).

Endurance training also increases the hepatic production of heat shock proteins and decreases the secretion of orixogenic proteins. Thus, endurance training in mice increased hepatic 70 kDa HSP (Mikami et al. 2004) and HSP72 (Atalay et al. 2004) expression in both the liver and other tissues, and the expression of hepatic orixogenic Agouti-related protein was reduced in rats after training (Ghanbari-Niaki et al. 2009); the latter change likely reduces the animals' appetite.

#### Triggers of hepatic responses

Both changes in the concentrations of hormones (insulin, glucagon and oestrogen) and cytokines (IL-1 $\beta$ , IL-6, IL-10 and IGF-1) and altered tissue sensitivity to these agents may contribute to the changes of hepatic metabolism observed following aerobic training.

Hepatic insulin sensitivity was increased in some animal studies. Regular exercise reduced the hepatic mRNA level and protein content of hepatic PEPCK, thus contributing to the improved insulin sensitivity (Chang et al. 2006).

However, training has not enhanced hepatic insulin sensitivity in all human studies (Hickman et al. 2004).

A week of repeated bouts of swimming reduced the liver fat content of male rats (Hu et al. 2000; Peijie et al. 2004). Regular training also increased the hepatic glucagon receptor density and glucocorticoid receptor count in exercise-trained rats (Légaré et al. 2001); an increased availability of glucagon may be important to this effect of exercise training, since liver fat was not reduced in animals that lacked glucagon receptors (Berglund et al. 2011).

Hepatic oestrogen receptors appear to influence the effects of exercise training on hepatic lipid metabolism (Paquette et al. 2007). Ovariectomy predisposes rats to hepatic steatosis, an increase of inflammatory biomarkers (e.g. inhibitor- $\kappa$ B kinase  $\beta$  and interleukin-6), an increased activity of hepatic lipogenic enzymes (e.g. sterol regulatory element-binding protein-1c, acetyl-CoA carboxylase (ACC) and stearoyl CoA desaturase), and a decreased expression of enzymes related to fat oxidation (e.g. carnitine palmitoyltransferase and hydroxyacyl-CoA-dehydrogenase). With the exception of increases in ACC, these adverse changes could be reversed, at least in rats and mice, through regular exercise (Jackson et al. 2011; Pighon et al. 2011; Domingos et al. 2012). Carnitine is an important cofactor for the oxidation of both long-chained fatty acids and carbohydrate, and may itself play an important role in the hepatic response; regular exercise attenuates the high-fat diet-induced reduction in carnitine palmitoyltransferase I activity (Cha et al. 2003), and up-regulates the genes involved in hepatic carnitine synthesis and uptake (Ringseis et al. 2011). Training also increased gene expression of microsomal triglyceride transfer protein and diacylglycerol acyltransferase-2 in ovariectomized rats, with a reduction in hepatic triglyceride content (Barsalani et al. 2010).

Some studies have observed greater serum levels of free IGF following aerobic training, either via increased hepatic IGF production (Prior et al. 2012), or because of increased hydrolysis of the corresponding binding factor (Schwarz et al. 1996). Resistance training likely has a similar effect (Bermon et al. 1999). On the other hand, a combination of regular exercise and a low fat diet increased serum concentrations of IGF-1 binding protein, thus decreasing circulating levels of free IGF-1, both in rats and in humans (Nishida et al. 2010; Wiczorek-Baranowska et al. 2011).

Rodent studies have suggested that regular aerobic exercise training may decrease tissue levels of the inflammatory cytokines IL-6 (Moon et al. 2012b) and IL-1 $\beta$  (de Araújo et al. 2012), and increase levels of the anti-inflammatory cytokine IL-10 (with an associated decrease in hepatic apoptosis) (de Araújo et al. 2012). Whilst there is some evidence from human investigations demonstrating a net hepatosplanchnic uptake of IL-6 during moderate intensity exercise (Febbraio et al. 2003), it remains to be determined whether

the liver is merely clearing this cytokine from the circulation, or whether it has a specific role in glucose homeostasis.

In conclusion, a variety of triggers have been suggested for the adaptations of hepatic metabolism associated with exercise training, including alterations in concentrations of and sensitivity to hormones (glucagon, insulin, oestrogen and IGF) and cytokines (IL-1 $\beta$ , IL-6 and IL-10); further research is needed to determine which are important factors, and which are incidental consequences of the observed adaptations.

#### Role of oxidant stress

Oxidant stress reflects a disequilibrium between the protein load and the ability of the hepatocyte endoplasmic reticulum to fold and assemble proteins correctly. It can be caused either by aging or by severe exercise, with an increased production of superoxides, a decrease of buffering agents, and/or a decrease of peroxidases. In rats, prolonged bouts of vigorous exercise (2 h swimming/day for 3 months) led to a down-regulation of cytosolic aconitase, a key factor in cellular iron homeostasis (Ho et al. 2001), possibly because of an increased production of NO and oxidative stress.

Most studies of mice, rats and dogs (Table 4) have shown moderate aerobic training as minimizing oxidative stress. Markers of oxidative stress are decreased (Navarro et al. 2003), and the activities of hepatic antioxidant enzymes such as superoxide dismutase (Gore et al. 1998; Burneiko et al. 2006; da Silva et al. 2009) and the corresponding signaling molecules (Huang et al. 2010) are increased. Further, hepatic glutathione transferase S activity and concentrations of reduced glutathione are increased (Sen et al. 1992; Radak et al. 2004), and the gene expression of unfolded protein response markers is enhanced (Chapados and Lavoie 2010).

Nevertheless, a few investigators have found no change or even a decrease of anti-oxidant enzyme activity following heavy endurance training (Hong and Johnson 1995), with a decreased hepatic superoxide dismutase mRNA, but an increase of catalase mRNA (Wilson and Johnson 2000), and [contrary to early observations on isolated hepatocytes (Eklöv et al. 1984)], little relationship between anti-oxidant enzyme levels and local oxidant stress (Ji et al. 1990; Godin and Garnett 1992).

Thus, we may conclude that moderate exercise training reduces hepatic oxidant stress, but very heavy training may have adverse effects upon oxidant status.

#### Functional activity

It is unclear from human studies of serum enzyme levels and pharmacokinetics how far liver function is influenced



either by habitual physical activity or by regular low to moderate intensities of exercise training. Nevertheless, the traditional clinical markers of hepatic function (serum ALT and GGT levels) do show a negative correlation with habitual physical activity (Robinson and Whitehead 1989; Nilsen et al. 1990; Pintus and Mascia 1996), probably because a sedentary lifestyle predisposes to steatosis (Whitfield 2001). Whether there is a more direct relationship between physical activity level, fitness and serum aminotransferase levels is discussed below.

In terms of pharmacokinetics, exercise training did not alter creatinine clearance in boxers (Saengsirisuwan et al. 1998), or the pharmacodynamics of propranolol in sedentary subjects (Frank et al. 1990; Panton et al. 1995). Likewise, cross-sectional research showed no significant differences in aminopyrine metabolism, galactose elimination, or indocyanine green clearance between endurance runners and relatively sedentary medical students (Ducry et al. 1979).

However, other reports suggest that hepatic function may be enhanced by vigorous (but not exhausting) training (Døssing 1985). Thus, the clearance of antipyrine (which depends almost exclusively upon hepatic metabolism) was faster in athletes than in controls, with no difference between sprinters and endurance competitors (Orioli et al. 1990). Likewise, endurance runners had a faster clearance of antipyrine than sedentary but otherwise healthy men (Villa et al. 1998). Longitudinal evidence supports these cross-sectional inferences. Three months of exercise training increased the clearance of antipyrine and aminopyrine in previously inactive students; moreover, individual improvements in these indices correlated highly with gains in  $\dot{V}O_{2max}$ , which averaged 6 % (Boel et al. 1984). Three months of moderate intensity exercise (combined aerobic and resistance training) also increased antipyrine clearance in elderly women (Mauriz et al. 2000).

Animal experiments generally confirm the beneficial effects of regular exercise on liver function seen in human subjects. Five weeks of training increased antipyrine clearance in mares (Dyke et al. 1998), and the livers of regularly exercised rats had a greater ability to metabolize and excrete certain chemicals not normally found in the body, such as naphthol and styrene products (Yiamouyiannis et al. 1992) and halothane (Daggan et al. 2000). In the study of halothane toxicity, hepatic glutathione levels were unchanged by 10 weeks of treadmill exercise, and it remained unclear whether benefit was due to enhanced antioxidant defence mechanisms or the associated decrease in hepatic fat (Daggan et al. 2000).

A further factor increasing the liver's ability to eliminate some substances is an increased secretion of biliary transporters. Chronic exercise such as swimming or running augments the hepatic production of bile acids (Frenkl et al. 1980) and increases the availability of bile acid transporters

(Yiamouyiannis et al. 1993). These changes may accelerate biliary clearance (but not necessarily blood stream clearance) of substances such as indocyanine green (Yiamouyiannis et al. 1993) acetaminophen and antipyrine (Frenkl et al. 1980).

Thus, the general impression from studies of pharmacokinetics is that regular moderate exercise enhances the functional clearance capacity of the liver.

### Summary of responses to chronic exercise training

Regular exercise training increases liver glycogen storage and the hepatic capacity for glucose output. On the other hand, glycogenolysis and gluconeogenesis are reduced at a given work-rate after training, with a reduced availability of gluconogenic precursors. The net effect is an improved ability to maintain euglycaemia, probably triggered by changes in hormone concentrations and sensitivity. Regular exercise training appears to reduce overall liver mass and associated fat mass, with an increase in HDL-c levels. Hepatic albumin and HSPs increase, and orixogenic proteins decrease with regular exercise training. The precise triggers for these changes of hepatic function are contentious, although hormones (insulin, glucagon and oestrogen) and a number of cytokines appear to be involved. Most (but not all) studies suggest that regular exercise training reduces markers of oxidative stress and increases antioxidant enzyme levels in the liver. Evidence for the overall effect of exercise training in terms of serum aminotransferase levels is conflicting, but most cross-sectional and longitudinal research indicates an improvement of hepatic clearance function with regular exercise.

### The role of physical activity in liver disease

In the final section of this review, we will examine interactions between physical activity and certain chronic liver conditions, including non-alcoholic fatty liver disease (NAFLD), hepatic inflammation and cirrhosis, and hepatic carcinoma, considering specifically the roles of inadequate habitual physical activity and co-pathologies in the genesis of these syndromes. We will also examine the impact of exercise training upon liver fat, as seen in both cross-sectional and longitudinal studies, and will finally consider appropriate exercise dose recommendations for the treatment of these disorders.

#### Non-alcoholic fatty liver disease (NAFLD)

NAFLD is characterized by the accumulation of fat in hepatocytes in the absence of excessive alcohol

consumption. The liver normally contains some fat (the triglycerides stored in hepatocytes), but NAFLD is commonly diagnosed when fat stores exceed ~5 % of hepatic mass. NAFLD accounts for the majority of liver disease worldwide; the condition is thought to affect up to one-third of adults (Browning et al. 2004; Szczepaniak et al. 2005) and it is found in most individuals who are obese (Bellentani et al. 2000). Even in children, the prevalence of NAFLD ranges from 2.6 to 9.6 %, depending upon age, sex, ethnic group and habitual physical activity (Takahashi and Fukusato 2010; Tsuruta et al. 2010). Liver biopsy and histological assessment provides the gold standard for the diagnosis of NAFLD, but in human research studies the liver fat content has more commonly been inferred from proton magnetic resonance spectroscopy or CT scan, and in animal experimentation the usual approach has been tissue analysis at sacrifice.

Hepatic fat accumulation is commonly associated with obesity, cardiovascular disease and diabetes. The build-up of triglycerides in the liver could reflect an increased delivery of fatty acids either from adipose tissue or directly from the diet, increased *de novo* hepatic lipogenesis, decreased hepatic fatty acid oxidation, or a decreased exit of fatty acids from the liver. The first of these mechanisms is probably the most important (Katsanos 2004); it accounts for the major fraction of fatty acids incorporated into liver fat in obese individuals under fasting conditions (Donnelly et al. 2005). The increase in hepatic fat may impair the insulin sensitivity of the hepatocytes (above), and insulin resistance is also manifest in adipose tissue (Kotronen et al. 2008), so that any given secretion of insulin is less effective in reducing lipolysis (Korenblat et al. 2008).

NAFLD can progress from a simple accumulation of fat through inflammation (steato-hepatitis) to fibrosis, cirrhosis and liver failure and even hepatic carcinoma (Angulo 2002). It is not entirely clear why the condition remains a simple steatosis in some individuals, but shows a progression of pathology in others. Inter-individual differences in reactions to reactive oxygen species, cytotoxic dicarboxylic acids, and hormonal balance, as well as mitochondrial abnormalities may be involved (Angulo 2002). Progression from simple steatosis to steatohepatitis probably reflects the combined effects of hepatic fat accumulation and oxidative stress, possibly exacerbated by endoplasmic reticulum stress (Malhi and Kaufman 2011) and gut barrier dysfunction (Rao 2009); anti-oxidant therapy is not necessarily helpful in preventing disease progression (Nobili et al. 2008).

#### Hepatic inflammation and cirrhosis

There is relatively little research evidence concerning interactions between physical activity and the more advanced

stages in the spectrum of NAFLD. From available information, it could be suggested that physical activity has some direct positive influence on hepatic pathology beyond simply modifying liver fat levels. For instance, as fibrosis develops, markers of hepatic apoptosis [plasma cytokeratin 18 (CK18) fragments, soluble Fas (sFas), and sFas ligand (sFasL)] increase (Fealy et al. 2012), and these changes have been positively associated with physical inactivity (Lee et al. 2008).

The situation can become a vicious cycle, since any form of hepatitis may discourage physical activity. Individuals affected by chronic hepatitis C infection were found to be less active than their peers, and to engage in less vigorous activity (Moon et al. 2012a). The intensity of physical activity seems important in preventing disease progression, since in a large adult cohort with biopsy-proven steatosis, neither total reported exercise per week nor the duration of moderate physical activity was associated with either the risk of steatohepatitis or the histological stage of fibrosis. On the other hand, meeting the weekly vigorous physical activity recommendation reduced the odds of steato-hepatitis to 0.65, and spending double the recommended time in vigorous activity also reduced the odds of advanced fibrosis to 0.53 (Kistler et al. 2011).

Three months of moderate intensity exercise therapy (5 days/week) lowered serum amino-transferases (ALT and AST) in patients with cirrhosis (Baba et al. 2006), and as little as a week of vigorous exercise training was sufficient to decrease ALT and CK-18 fragments (Fealy et al. 2012). Nevertheless, the primary rationale for advocating exercise therapy in patients with advanced liver disease is arguably for the multiplicity of other benefits of chronic exercise, especially those relating to physical weakness and comorbidities. Thirteen studies of patients with hepatic cirrhosis noted substantial decreases in aerobic capacity and muscular strength relative to healthy controls (Jones et al. 2012). Low levels of aerobic fitness and exercise tolerance (Wiesinger et al. 2001; Pieber et al. 2006; Dharancy et al. 2008) have been confirmed in other studies (Ritland et al. 1982, 1983; Campillo et al. 1990a, b; DeLissio et al. 1991; Terziyski et al. 2008), particularly in individuals with associated ascites (Campillo et al. 1990b; Wong et al. 2001). There is also evidence of muscular weakness (Tarter et al. 1997; Andersen et al. 1998), proportional to the severity of disease, but independent of its etiology (Campillo et al. 1990b; Wiesinger et al. 2001; Terziyski et al. 2008).

Thus, exercise that includes an element of resistance training is arguably a useful therapy for improving fitness and functional capacity in this population, but it remains unclear whether exercise can restore liver health (and if so, the dose that is needed). One major obstacle to implementing and sustaining a programme of exercise training in advanced liver disease is initial fatigue; this has an adverse effect upon the

individual's quality of life (Stanca et al. 2005), and by discouraging physical activity, it progressively exacerbates the initial loss of muscular strength (Wu et al. 2012). Nevertheless, a programme of regular progressive exercise can counter fatigue, even in people with advanced fibrosis (Zucker 2004). Moreover, given adequate motivation, patients with cirrhosis can tolerate quite vigorous exercise, maintaining oxygenation of the brain and muscles (Bay Nielsen et al. 2005) and showing no evidence of hypoglycemia while they are active (DeLissio et al. 1991). The one major concern in this condition is the potential to develop oesophageal bleeding. The hepatic venous pressure gradient is increased even at low levels of physical activity (30 % of peak work-rate) (Garcia-Pagan et al. 1996), and in patients with oesophageal varices, portal hypertension induced by over-vigorous exercise could cause such bleeding.

There have been few investigations of the effect of aerobic training in hepatic cirrhosis. One investigation reported a 29 % gain of predicted  $\dot{V}O_{2max}$  over 10–12 weeks of training (Ritland et al. 1983), and a second trial with only four subjects found an increase of  $\dot{V}O_{2max}$  in two of the four individuals, with an 18–20 % improvement of muscle strength in these two individuals (Campillo et al. 1990b).

#### Animal studies of exercise and liver pathologies

Animal studies have underlined the potential of exercise to have direct beneficial effects upon the diseased liver. In one such study, mice were fed a high-fat diet; however, those animals that subsequently underwent a progressive 16-week aerobic exercise intervention showed an elevation of hepatic tumor necrosis factor levels, together with a reduction or abolition of macrophage infiltration and signs of fibrosis (Sirius red and -smooth muscle actin staining and tissue inhibitor of matrix metalloproteinase-1 mRNA) (Kawanishi et al. 2012). A second investigation noted signs of inflammation and steatohepatitis (macro-vesicular steatosis and lymphocytosis) in sedentary rats that were fed a high-fat diet, but such findings were greatly attenuated in their peers who exercised daily; ALT but not AST levels were also reduced in exercised animals (He et al. 2008).

We may thus conclude that exercise programmes have favourable effects in advanced hepatic disease, provided that patients can be motivated to sustain such activity.

#### Hepatocellular carcinoma

There has been a paucity of research into interactions between physical activity and hepato-cellular carcinoma. A 10-year follow-up of study of 507,897 retired Americans found a significantly reduced risk of hepatic carcinoma in those who were regularly active (>5 times a week) vs. those who reported exercising never or rarely (odds ratio 0.64)

(Behrens et al. 2013). A moderate exercise programme may be beneficial, even if the hepatic carcinoma is quite advanced. One case report noted an increase in aerobic capacity after 6 weeks of supervised aerobic exercise therapy (Crevenna et al. 2003).

#### The influence of other pathologies associated with inadequate habitual physical activity

The vast majority of research concerning physical activity in the aetiology and management of liver disease has focused on simple hepatic steatosis (detailed below). However, NAFLD is commonly associated with other markers of inadequate habitual physical activity, including cardiovascular disease, metabolic syndrome and type 2 diabetes mellitus. In terms of associated insulin sensitivity, univariate correlations suggest that although body fatness is a prime determinant of whole-body insulin sensitivity, the main determinant of hepatic insulin sensitivity may be the individual's active energy expenditure (Holt et al. 2007). A follow-up of 6,003 patients with non-alcoholic fatty liver disease found 411 developed type 2 diabetes over a follow-up averaging 4.9 years; a Cox proportional hazards analysis demonstrated that a gamma glutamyl transferase (GGT) >109 IU/L and an exercise level of less than 60 min per week were significant predictors of diabetes, both with hazard ratios averaging 1.60 (Arase et al. 2009). GGT facilitates the intracellular transport of glutathione, and increases in levels of this enzyme are a possible indicator of oxidative stress, which in turn can predispose to diabetes (Nannipieri et al. 2005).

Whilst it has been thoroughly documented that low levels of habitual physical activity predispose to the obesity, dyslipidaemia, impaired glucose tolerance and high blood pressure that characterize cardiovascular disease, the metabolic syndrome and diabetes, and that a physical activity intervention is effective in their management (Winnick et al. 2008), fat reduction in the liver is also an important component of both prevention and treatment. A decrease of hepatic fat content has been thought to avert type 2 diabetes mellitus, particularly in older individuals (Tamura et al. 2005; Thamer et al. 2007). Similarly, a normalizing of liver fat content in patients with type 2 diabetes improves the insulin-induced suppression of hepatic glucose output and restores normal fasting blood glucose concentration (Petersen et al. 2005). Recent research interest has thus centred on the role of NAFLD in these pathologies, and the effect of physical activity on liver fat levels.

#### The association between habitual physical activity/fitness and liver fat (cross-sectional studies)

A possible role for exercise therapy in the management of NAFLD is supported by many cross-sectional

investigations that show an association between low levels of habitual physical activity and/or fitness and the prevalence of NAFLD. Sixteen such studies of human subjects have made cross-sectional assessments of habitual physical activity (Table 5); 12 used physical activity questionnaires, three used objective activity monitors (Newton et al. 2008; Fintini et al. 2012; Gerber et al. 2012), and one classified subjects based upon their obesity (Viitasalo et al. 2012). Sample size ranged from small groups to populations >30,000, and one analysis was based upon twins with dissimilar activity patterns (Leskinen et al. 2009). In one instance, objective monitoring suggested an effect of physical activity, but (probably because of lesser reliability and validity) subjective questionnaires completed by the same individuals did not (Fintini et al. 2012). Collectively, these studies showed that habitual physical activity was an important correlate of hepatic fat in most comparisons, with a possible exercise volume-response relationship (Hsieh et al. 1998), although two reports found no relationship between the severity of histological abnormalities and physical activity (Kang et al. 2006; Kistler et al. 2011).

Eleven reports [including the one twin study (Leskinen et al. 2009)] related cycle ergometer or treadmill assessments of aerobic fitness to hepatic fat accumulation (Table 6). With two exceptions (Seppala-Lindroos et al. 2007; Krasnoff et al. 2008), an inverse relationship was seen. However, in some studies the negative association was relatively weak (Nguyen-Duy et al. 2003; McMillan et al. 2007), particularly if data were co-varied for inter-individual differences in obesity.

#### Physical activity interventions and liver fat (longitudinal trials in humans)

Forty-six longitudinal human trials were identified (Table 7). Often, sample sizes were small; 19 trials included some form of non-exercise control group, often “usual treatment” or a dietary regimen. Interventions ranged from general lifestyle recommendations to specific programmes with careful control of both exercise and diet. Programmes typically yielded consistent reductions in liver fat, and this was usually associated with decreased insulin resistance. One report noted an improvement of histopathology in response to a combined exercise and weight loss programme (Goodpaster et al. 2010), but there is little evidence in this regard. Effects on serum aminotransferase levels have also been unclear, possibly confounded by normal or near-normal levels in the studied cohorts prior to interventions (Keating et al. 2012; Thoma et al. 2012).

It seems likely that benefits such as a reduction of hepatic fat and a possible normalization of serum aminotransferases will be maximized by a combination of physical activity and dieting which results in significant

weight loss, but the respective contributions of diet, physical activity and weight loss to improvements in hepatic function remain to be defined (Thoma et al. 2012). Exercise has traditionally been employed with the goal of weight loss, but some investigators have found benefits from exercise in the absence of dieting (Larson-Meyer et al. 2008) or any change in body mass (Johnson et al. 2009). Further, benefits have persisted after statistical adjustment of data for changes of body mass (Bonekamp et al. 2008). Moreover, at least one study found that dietary manipulation did not enhance the effects of exercise (Eckard et al. 2013).

Nevertheless, much of the current evidence suggests that exercise training may, at best, enhance the hepatic effects of dieting (Goodpaster et al. 2010), and may (Coker et al. 2009) or may not (Tamura et al. 2005; Shah et al. 2009) further increase the insulin sensitization induced by dieting. Significant weight loss (10 %) seems the most effective means to lower liver fat content and aminotransferase levels; lesser effects are seen in studies where the decrease in body mass was 5 % or less (Chen et al. 2008), or if exercise did not induce weight loss. Several reports have found that although exercise has other benefits, such as insulin sensitization, it does not enhance the hepatic response to dieting (Tamura et al. 2005; Shojaee-Moradie et al. 2007; Shah et al. 2009; van der Heijden et al. 2009, 2010b; Jenkins and Hagberg 2011; Straznicky et al. 2012).

Most investigations have evaluated aerobic training programmes. A few reports have also noted favourable responses to resistance exercise training, although its effectiveness in NAFLD is less clearly established. Two of three comparisons between aerobic and resistance training (Lee et al. 2012; Bacchi and Moghetti 2013) found similar decreases of hepatic fat with both types of exercise. However, the third and largest study found no benefit from resistance training alone, and the response to aerobic training was not enhanced by adding resistance activity (Slentz et al. 2012). Another study of a resistance exercise programme found no reduction of inflammatory markers (Levinger et al. 2009), and one 12-week trial of resistance exercise found a decrease of insulin resistance without a change of hepatic fat content (van der Heijden et al. 2010a). In contrast, a controlled 3-month trial in obese adolescent boys found that thrice weekly 60-min sessions of either aerobic exercise or resistance exercise reduced liver fat, but only resistance exercise was effective in increasing insulin sensitivity (Lee et al. 2012). It is plainly as yet unclear and important to resolve how effective resistance training is for decreasing steatosis and associated comorbidities, particularly as it has been suggested that resistance exercise is important to correct the muscular weakness and autonomic dysfunction that is often associated with this condition (Jakovljevic et al. 2013).

**Table 5** Cross-sectional analysis of the inverse association between reported physical activity and steatosis

Author	Sample	Liver assessment	Activity assessment	Outcome	Comment
Bae et al. (2012)	72,359 Korean adults without diabetes	Ultrasound, liver enzymes	Self-reported physical activity	Those exercising at least 3 times per week for at least 30 min per session over a period of 3 months had reduced risk of steatosis (0.53–0.72)	Active individuals also had a decreased risk of elevated hepatic AST (0.85) and ALT (0.74)
Fintini et al. (2012)	Comparison of children with steatosis ( $n = 40$ ) obese ( $n = 30$ ) and lean peers ( $n = 41$ )		Physical activity questionnaires and sense-wear arm bands	Sense-wear data showed steatosis group devoted more time to sedentary pursuits than lean, and engaged in less physical activity	Questionnaire data did not agree with Sense-wear data
Gerber et al. (2012)	3,056 participants in the US NHANES survey of 2003–2006 aged >20 years	Fatty liver index based on BMI, waist circumference, triglycerides and GGT	Accelerometer step counts	Individuals with high fatty liver index (>60 units) had lower levels of accelerometer readings (29 counts/min per day), and spent less time than controls at all levels of activity	
Hattar et al. (2011)	57 Hispanic children, average age 12.1 years (20 steatosis, 20 obese, 17 controls)	Liver biopsy, liver enzymes	Retrospective physical activity scores, based on questionnaire	Sedentary score >2 associated with stage 2–3 hepatic fibrosis	15 % of children with steatohepatitis performed light exercise, compared with 35 % of obese and 59 % of non-obese children
Hsieh et al. (1998)	3,331 adult Japanese men	Ultrasound	Physical activity questionnaire	Fatty liver less prevalent in those regularly exercising >2 days/week than in sedentary men	Dose–response relationship (sedentary vs. those active 1, 2 and 3 days/week)
Kang et al. (2006)	39 M, 52 F with steatosis, age 48 years	Liver biopsy	Physical activity questionnaire	No difference of histological severity with reported physical activity	
Kistler et al. (2011)	302 men, 511 women with steatosis	Histology	Self-reported physical activity	Neither moderate nor total exercise associated with stage of fibrosis; however, meeting vigorous recommendations reduced odds of steatohepatitis	
Lawlor et al. (2005)	3,789 British women aged 60–79 years	Liver enzymes (ALT, GGT)	British regional heart activity questionnaire	Frequency of activity inversely related to GGT, but relationship to ALT attenuated by adjustment for body mass index and waist hip ratio	
Lee et al. (2006)	201 children >140 % of ideal body mass	Liver enzymes	Physical activity questionnaire	53/201 children show increased ALT and AST; odds ratio if low activity 2.39	Increase of ALT and AST not well correlated with body mass or body fat, but associated with insulin resistance
Leskinen et al. (2009)	16 same-sex middle-aged twin pairs discordant for activity	Proton magnetic resonance spectroscopy	Physical activity questionnaire	Inactive twins had 3 times as much liver fat	Inactive twins also had greater body mass, fat mass, visceral fat and lower fitness
Mager et al. (2010)	38 children (36 obese) aged 5–19 years	Ultrasound and transferases	3-day reported dietary intake and activity questionnaire	Physical activity low in almost all children	No clear comparison group

Table 5 continued

Author	Sample	Liver assessment	Activity assessment	Outcome	Comment
Newton et al. (2008)	36 M, 36 F (36 with steatosis, 36 controls)	Liver biopsy	Pedometer (6-day record)	Cases of steatosis took 20 % fewer steps/day	
Perseghin et al. (2007)	114 M, 77 F aged 19–62 years	Proton magnetic resonance spectroscopy, Fatty liver defined as >5 % wet weight	Physical activity questionnaire	Prevalence of steatosis (defined as hepatic fat >5 %) 25 % in least active quartile, 2 % in most active quartile	Association attenuated by adjustments for age, sex, BMI, insulin sensitivity and adiponectin
Suzuki et al. (2005)	348 men with elevated ALT, mean age 42 year	ALT levels	Self-report	Regular exercise associated with lower ALT, and 2.5-fold greater likelihood of normalizing even after adjusting for change in body weight	
Tiikkainen et al. (2002)	27 women with previous gestational diabetes	Proton magnetic spectroscopy; categorized as hepatic fat above or below 5 %	Frequency of 30 min bouts of physical activity per week	Women with low fat exercising 5 times/week, those with steatosis only 3 times/week	Steatosis associated with insulin resistance
Vitasalo et al. (2012)	492 children aged 6–8 years	Hepatic enzymes	Activity not reported	Overweight or obesity increases risk of high ALT 2.1-fold, high GGT 4.5-fold	GGT linked to c-reactive protein in obesity, suggesting low-grade inflammation
Zelber-Sagi et al. (2008)	375 Israeli men and women, mean age 51 years	Abdominal ultra-sound	Self-reported physical activity	Hepatic fat related to sports participation (odds ratio 0.66) and resistance exercise (0.61), but only related to reports of resistance exercise after adjustment for BMI; even this relationship non-significant if also adjusted for leptin or waist circumference	

## Physical activity interventions and liver fat (longitudinal trials in animals)

Some 21 animal studies of exercise and hepatic steatosis generally confirm the findings of human longitudinal investigations (Table 8). They provide growing empirical evidence that fat accumulation has direct adverse effects upon hepatic function, and that these changes can be reversed by exercise; further, they add helpful information on cellular mechanisms underlying the adverse effect of hepatic fat upon glucose homeostasis (Table 9).

In mice fed a high-fat diet, regular exercise reduced the accumulation of fat in the liver, improved insulin resistance and reduced circulating cholesterol, triglycerides, and AST and ALT levels (Marques et al. 2010). Dietary restriction, voluntary wheel running and imposed swimming or treadmill running all seem effective in preventing steatosis (see Table 8 for references), and in one report hepatic benefits were elicited more readily by intermittent swim training than by continuous bouts of swimming (Sene-Fiorese et al. 2009). Yasari et al. (2006) found that after 6 weeks of detraining, rats trained on a treadmill for 4 weeks had regained a similar body fat to sedentary animals, although liver lipid infiltration was not increased with cessation of training. In contrast, Linden et al. (2013) found that 4 weeks of inactivity following 16 weeks of wheel running caused the development of hepatic steatosis in obese rats, although liver triglycerides were still 60 % lower than in animals that had remained sedentary throughout.

Among mechanisms underlying the adverse effect of hepatic fat upon glucose homeostasis, lipid accumulation appears to down-regulate phosphatidylinositol 3-kinase, an enzyme that has a central role in mediating the action of insulin in hepatocytes (Katsanos 2004). Rats fed an obesity-inducing diet not only developed peripheral insulin resistance, but also demonstrated endoplasmic reticular stress in both hepatic and adipose tissues, with activation of the proinflammatory molecules c-jun N-terminal kinase (JNK) and nuclear factor kappa-B (NF- $\kappa$ B).

Cellular adaptations associated with the benefits of enhanced activity have included increased hepatic mitochondrial fatty acid oxidation, enhanced oxidative enzyme function and protein content, and suppression of de novo lipogenesis (Rector et al. 2011). Specific molecular mechanisms identified as contributing to attenuation of fat accumulation and/or reversal of steatosis have included increased hepatic mitochondrial activity (citrate synthase,  $\beta$ -hydroxyacyl-dehydrogenase [HAD] and cytochrome c oxidase) and subsequent beta-oxidation (Rector et al. 2011), a decrease of regulatory element-binding protein-1c (SREBP-1c, one of a group of transcription factors regulating the genes involved in cholesterol and fatty acid synthesis) (Cintra et al. 2012), down-regulation of the hepatic

SCD-1 gene, and thus of SCD-1, a rate-limiting enzyme in the biosynthesis of monounsaturated fats (Yasari et al. 2010), and a decreased activity of the hepatic ketone synthesis pathway seen in streptozotocin-diabetic rats, with a decreased activity of the corresponding rate-limiting enzyme HMG-CoA (El Midaoui et al. 2006). Whereas streptozotocin diabetic rats showed a greatly increased activity of the branched-chain alpha-ketoacid dehydrogenase (BCKDH) complex, the rate-limiting enzyme in the catabolism of branched-chain amino acids, such activity was normalized by regular exercise (Li et al. 2001). Regular exercise also attenuated the reduction in hepatic IGF-1 seen in alloxan-diabetic rats (Leme et al. 2009).

Exercise training also reduced hepatic JNK and NF- $\kappa$ B, and lessened endoplasmic reticular stress as shown by decreasing phosphorylation of the two major metabolic markers of this condition (protein-kinase like endoplasmic reticular kinase, PERK and eukaryotic initiation factor 2, eIF2 phosphorylation) (da Luz et al. 2011). Moreover, the glucose stimulation of insulin secretion was decreased in rats that were given access to an exercise wheel, without any deterioration in glucose homeostasis; activity of the insulin-inducible enzyme hepatic glucose kinase (the first stage in glucose utilization) was decreased, possibly due to the lesser output of insulin (Zawalich et al. 1982). Seven days of voluntary wheel running increased the release of the hormone-like hepatic insulin sensitizing substance (HISS), thus decreasing the peripheral insulin resistance of rats (Chowdhury et al. 2013). Aging decreases the hepatic output of HISS, but again this could be countered by allowing the rat free access to a running wheel (Chowdhury et al. 2011). Exercise partially reversed attenuated insulin and leptin signalling in chlorpromazine-induced diabetes in rats by increasing concentrations of insulin-receptor substrate-2 protein (Park et al. 2007). In exercised mice, the enhanced insulin sensitivity was associated with an increased hepatic expression of endosomal adaptor protein APPL1, which blocks the association of protein kinase AKT with its endogenous inhibitor tribbles-related protein 3 (TRB3), and there was a decreased expression of TRB3 (Marinho et al. 2012).

In contrast, several metabolic precursors of steatosis were seen in hyperphagic obese rats following a sudden 1-week cessation of exercise. Changes included a decrease in hepatic mitochondrial oxidative capacity, an increased hepatic expression of lipogenic proteins, and increased levels of hepatic malonyl CoA (Rector et al. 2008).

Additional effects of exercise training upon insulin sensitivity arise outside the liver, from an increase in muscle mass, an alteration in muscle quality, the greater energy demands of skeletal muscle, and the reduction of visceral fat stores (with a lesser incorporation of fatty acids into the liver). Exercise programmes may also influence

**Table 6** Cross-sectional analysis of benefits of cardiorespiratory fitness in steatosis

Author	Sample	Liver assessment	Fitness assessment	Outcome	Comment
Church et al. (2006)	218 men aged 33–73 years	CT scan, ALT > 30 U/L, AST/ALT ratio <1.0	Peak treadmill endurance time	Treadmill endurance and BMI independently associated with steatosis	Associations attenuated if abdominal fatness included in model
Hannukainen et al. (2007)	Nine male monozygotic twin pairs differing in habitual activity	Proton magnetic resonance spectroscopy	Cycle ergometry (active twins had 18 % higher maximal oxygen intake)	20 % less visceral fat in more active twins	Hepatic uptake of free fatty acids lower in active twins
Haufe et al. (2010)	Overweight and obese subjects (31 M, 108 F aged 40–50 years)	Proton magnetic resonance spectroscopy	Cycle ergometry, peak oxygen intake	Negative correlation between aerobic fitness and hepatic fat, M > F	In men, correlation independent of visceral adipose tissue
Kantartzis et al. (2009)	70 M, 100 F (50 with steatosis)	Proton magnetic resonance spectroscopy	Peak oxygen intake on cycle ergometer	Initial peak oxygen intake strongest predictor of reduction in hepatic fat with diet and physical activity lifestyle programme	Authors conclude cardiorespiratory fitness determines liver fat content
Krasnoff et al. (2008)	19 M, 18 F, average age 45.9 years	Steatosis classified by biopsy	Symptom-limited peak oxygen intake on treadmill	No relationship between peak oxygen intake and steatosis, but lower aerobic power if hepatitis (34.0 vs. 25.1 ml/kg.min)	Histological severity of liver disease influences health-related fitness
Kuk et al. (2004)	86 lean premenopausal women	CT scan	Peak treadmill endurance time	Treadmill endurance lower in those with steatosis	Liver fat not related to other metabolic risk factors
Leskinen et al. (2009)	16 same-sex middle-aged twin pairs discordant for activity	Proton magnetic resonance spectroscopy	Peak oxygen intake on cycle ergometer	Inactive twins had 3 times as much liver fat	Inactive also had greater body mass, fat mass, visceral fat and lower fitness
McMillan et al. (2007)	293 men aged 29–78 years	CT scan	Peak treadmill endurance time	Peak treadmill time weakly correlated with liver fat ( $r = -0.24$ )	Difference of predicted peak oxygen intake small after control for adiposity (35.3 vs. 34.1 ml/kg.min)
Nguyen-Duy et al. (2003)	161 men aged 33–72 years	CT scan	Peak treadmill endurance time	Peak treadmill time weakly correlated with liver fat ( $r = -0.26$ )	Probable overlap of sample with McMillan et al. (2007)
O'Donovan et al. (2009)	50 men aged 34–56, both obese and lean	Proton magnetic resonance spectroscopy	Peak oxygen intake on cycle ergometer	Liver fat was greater in unfit	Relationship eliminated by introduction of waist circumference
Seppala-Lindroos et al. (2007)	30 middle-aged men (15 with steatosis)	Proton magnetic resonance spectroscopy	Directly measured maximal oxygen intake	No significant difference of maximal oxygen intake between high and low fat groups (35.6 vs. 33.5 ml/kg min)	High liver fat defined as >3 %, so that some normal individuals in supposed high fat group?



**Table 7** Longitudinal human studies of changes in hepatic fat content with exercise interventions

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Albu et al. (2010)	58 obese, average age 59 years	CT scan	175-min moderate aerobic exercise/week + moderate energy restriction	No control group	18 % decrease in hepatic fat content
Bacchi and Moghetti (2013); Bacchi et al. (2013)	31 overweight or obese	Magnetic resonance imaging	4-month programme; 60 min aerobic exercise at 60–65 % heart rate reserve 3 times/week vs. 60 min resistance exercise at 70–80 % IRM 3 times/week	Diet unchanged. No non-exercise control	Hepatic fat reduced 33 % (aerobic) vs. 26 % resistance programme
Barsalani et al. (2013)	54 overweight or obese post-menopausal women	Liver enzymes and liver index	6 months of exercise (3 times/week, mixed training) + placebo, or similar exercise + flavones	No non-exercise control	Fatty index reduced 29 % in combined treatment, 18 % with exercise alone
Bonekamp et al. (2008)	45 adults with type 2 diabetes mellitus, mean age 53 years	Proton magnetic resonance spectroscopy	45 min of moderate aerobic exercise plus weight lifting, 3 times/week for 6 months, vs. control group	Dietary policy unclear	Significant reduction in hepatic fat (2.5 % absolute change), effect persisted even if adjusted for BMI or visceral fat. Abstract report only
Bozzetto et al. (2012)	45 obese diabetic subjects aged 35–70 years	Proton magnetic resonance spectroscopy	8-week aerobic exercise, 45 min at 70 % of peak oxygen intake, 2 times/week	4-way trial, with 2 diets (CHO/fibre or multiunsaturated fat)	Multi-unsaturated fat reduced liver fat 29 %; no added effect from exercise
Chen et al. (2008)	54 M and F Taiwanese with steatosis, average age 38–40 years	Ultrasound. Liver enzymes (AST, ALT, GGT)	10-week diet + exercise (high-intensity cycle ergometry, 1 h twice/week) vs. exercise alone vs. control	No non-exercise control	Diet + exercise decreased liver fat. ALT and GGT; exercise alone less effective, only decreased liver fat
de Piano et al. (2012)	58 post-pubertal obese adolescents, with or without steatosis, average age 16.5 years	Ultrasound, ALT	1-year lifestyle intervention with aerobic (60 min at ventilatory threshold, 3 times/week) or aerobic + resistance (3 sets of 6–20 reps for main muscle groups) training	No non-exercise control	Combined aerobic + resistance therapy more effective in improving ALT, insulin resistance and non-invasive markers of steatosis than aerobic exercise alone
Devries et al. (2008)	41 men and women, aged 38–40 years; half of sample lean, half obese	CT scan, liver enzymes	3 months of cycle ergometer training, 60 min/day, 3 times/week, progressing to 65 % of maximal oxygen intake in women and 70 % in men	Liver attenuation negatively correlated with ALT. No control group	Training did not alter liver attenuation on CT scan or ALT, but it reduced GGT in men only

Table 7 continued

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Eckard et al. (2013)	56 adults aged 18–70 with steatosis	Biopsy, liver enzymes	6-month programme; moderate exercise (30–60 min, 4–7 times/week, tracked by exercise log and pedometer) vs. exercise + low fat or moderate fat + low processed carbohydrates vs. standard care	All intervention groups improved, with no significant inter-group differences	
Fealy et al. (2012)	13 obese aged 58 years, sex not specified	Proton magnetic resonance spectroscopy	1 week of walking, 60 min/day at 85 % of maximal heart rate	Liver enzymes, CK18 markers of apoptosis. No control group	Reduced markers of apoptosis, mediated through increased oxidative capacity and greater insulin sensitivity
Finucane et al. (2010)	100 healthy older people (50 served as controls)	Proton magnetic resonance spectroscopy	12-week cycle ergometer exercise, 60 min, 3 times/week vs. control group	Increase of predicted maximal oxygen intake, no change of body mass	Significant reduction of liver fat in intervention group
Franzese et al. (1997)	41 M, 34 F obese children age 9.5 years	Ultrasound, liver enzymes	6 months of diet + exercise	No control group	Liver fat and transaminases decreased with loss of weight
Goodpaster et al. (2010)	130 severely obese adults (101 completed trial)	CT scan	6 months, diet + exercise vs. diet		Both groups lost weight, exercised group lost more liver fat
Grønbaek et al. (2012)	117 obese children, average age 12.1 years	Ultrasound	10-week period at a weight-loss camp with daily hour of varied aerobic exercise	Insulin sensitivity increased. No control group	Reduction of steatosis, decreased ALT in all participants; two-thirds of improvement in liver score retained at 12 months
Hallsworth et al. (2011)	19 sedentary adults with steatosis (average age 52 years experimental, 62 years controls)	Proton magnetic resonance spectroscopy	8-week resistance exercise ( $n = 11$ ) vs. standard treatment ( $n = 8$ ); physical activity monitored by sensor wear arm band	Lipid oxidation, glucose control and insulin resistance all improved. Control group older	13 % reduction of liver fat with no change in body mass, total fat mass or visceral fat volume
Hickman et al. (2004)	35 men and women (21 with hepatitis C virus), average age 44 years	Liver biopsy and enzymes	15-month programme of diet plus encouraging 150 min aerobic exercise per week	Exercise monitored by activity diaries. No control group	Improvement of ALT levels correlated with development and maintenance of weight loss; 14 patients biopsied after 3–6 months of exercise showed lessening of steatosis
Jin et al. (2012)	120 potential liver donors with steatosis	Hepatic biopsy	Dietary restriction + 10 weeks of exercise (three 20-min sessions of jogging or walking per week)	Improvement of steatosis associated with weight reduction >5 % and cholesterol reduction >10 %. No control group	Histological improvement in 103 of 120 subjects

**Table 7** continued

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Johnson et al. (2009)	19 sedentary obese men and women	Proton magnetic resonance spectroscopy	4 weeks of aerobic cycle ergometer exercise (30–45 min, 3 times/week) at intensity rising to 70 % of peak oxygen intake	No change of body mass. No control group	21 % reduction of hepatic triglycerides
Kawaguchi et al. (2011)	35 adults with steatosis resistant to lifestyle counselling (12 trained, 23 controls)	Ultrasound and liver enzymes	Hybrid training (voluntary and electrical contraction of quadriceps and hamstrings, 19 min, 2 times/week for 12 weeks	Associated reduction of insulin resistance and small decrease of body mass. No control group	Steatosis reduced, ALT decreased in trained group
Koot et al. (2011)	144 obese children, mean age 14.1 years	Ultrasound	6-month programme, with three 1-h sessions of unspecified exercise/week plus changes in eating behaviour	Changes related to decreased insulin resistance and 12 % decrease of body mass. No control group	Prevalence of steatosis decreased from 31 to 12 %. Prevalence of elevated ALT decreased from 26 to 11 %, elevated AST decreased from 13 to 4 % of sample
Kugelmas et al. (2003)	16 patients with steatosis, 18–65 years	Hepatic enzymes	12-week programme of diet and encouragement to walk or jog 30 min/day; half of group also received 800 IU vitamin E daily	No non-exercise control group	Reduction of ALT and AST in first 6 weeks, benefits sustained at 12 weeks
Larson-Meyer et al. (2008)	46 overweight men and women	Proton magnetic resonance spectroscopy, CT scan, liver enzymes	6-month study: dietary restriction + exercise (structured to increase energy expenditure 12.5 %) vs. dietary restriction vs. low-calorie diet vs. control	Liver fat initially correlated with ALT	Liver fat reduced in all three experimental groups; changes not associated with ALT or inflammatory markers
Lazo et al. (2008)	96 men and women with diabetes mellitus, aged 45–76 years, divided between lifestyle intervention and education/support group	Proton magnetic resonance spectroscopy	Weekly meetings to encourage dieting and progression to 175 min of moderate exercise per week for 1 year	Subjects instructed to follow weight maintenance diet; only resistance exercise increased insulin sensitivity	Decrease in steatosis 50.8 % (intervention) vs. 22.8 % (controls). No changes in AST or ALT
Lee et al. (2012)	48 obese adolescent boys divided into three groups	Proton magnetic resonance spectroscopy	3-month trial, 3 sessions/week 60 min aerobic exercise (50 % rising to 60–75 % of maximal oxygen intake) or resistance exercise (60 % of initial 1RM) or controls		Both types of exercise reduced hepatic lipids
Levinger et al. (2009)	28 M, 27 F aged 50.8 years (high vs. low metabolic risk factors)	Liver enzymes, inflammatory markers	10 weeks of resistance training 3 days/week, 50 % rising to 75–85 % 1RM exercise vs. controls	Risk factors associated with enzymes and inflammatory markers	Liver enzymes and inflammatory markers not reduced in those with high metabolic risk factors

Table 7 continued

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Nobili et al. (2008)	37 boys, 16 girls, aged 5.7–18.8 years (plus 33 drop-outs)	Liver enzymes and biopsy (lobular and portal inflammation and ballooning)	Diet plus physical activity for 24 months	Weight loss, insulin sensitivity, physical activity recall. No non-exercise control group	Improvement of enzymes and histology, not further improved by alpha tocopherol and ascorbic acid
Oza et al. (2009)	67 cases of steatosis, only 22 completed treatment	CT scan, liver enzymes	6-month home-based diet + exercise (target of 23 MET-h/week physical activity + 4 MET-h/week of exercise)	Poor compliance, no control group	19/22 showed decreases of visceral fat, liver/spleen ratio, AST, ALT, GGT
Promrat et al. (2010)	Overweight or obese subjects randomized to lifestyle (20) or education (10) groups	Histology	48-week intervention; weekly, then biweekly counselling aiming for 7–10 % weight loss, exercise and altered behaviour	Improvements in hepatic condition associated with weight loss	Histology improvement >3 points in 14/20 of intervention vs. 3/10 of control subjects
Rohrer et al. (2008)	16 boys, six girls, median age 11.9 years	Liver enzymes	1-year weight loss programme that included physical activity	Gains of physical strength and endurance. No control group	Improvement of hepatic enzymes associated with improved fitness
Santomauro et al. (2012)	36 obese children with steatosis	Ultrasound	1-year lifestyle approach, focusing on dieting and increased physical activity	Weight loss main variable accounting for reduction of steatosis. No control group	Reduction or disappearance of steatosis linked to increased physical activity and associated weight loss in 12 of 36 children
Schäfer et al. (2007)	48 impaired glucose tolerance, 133 normal subjects	Proton magnetic resonance spectroscopy	24-month diet + exercise at anaerobic threshold (Polar heart rate monitor)	Body mass decreased 3 % in both groups. No non-exercise control group	Liver fat decreased 28 %, visceral fat 8 % in those with impaired glucose tolerance
Shah et al. (2009)	18 obese >65 years	Proton magnetic resonance spectroscopy	Diet vs. diet + exercise (thrice weekly 90-min sessions of aerobic, resistance, flexibility and resistance training, with diet adjusted to achieve a similar energy deficit)	Body mass decreased 9–10 % in both groups. No non-exercise control	Both treatments reduced liver fat and insulin resistance to similar extent, but added exercise improved physical function
Shojaee-Moradie et al. (2007)	17 initially sedentary men (seven of 17 served as controls)	Proton magnetic resonance spectroscopy	Exercised for 6 weeks. 60–85 % of maximal aerobic power for 20 min, 3 times/week	No control group	No change of liver fat content, but decrease in hepatic insulin resistance and increase in maximal oxygen intake
Slentz et al. (2012)	155 overweight adults aged 18–70 years	Computed tomography, ALT	8-month training 3 days/week: aerobic (running 19 km/week at 75 % aerobic power) vs. resistance vs. combined aerobic + resistance training	No non-exercise controls. ALT inversely related to hepatic fat content	Aerobic training decreased hepatic fat 5.6 %. Also 13.7 % decrease of ALT. No benefit from resistance training alone. Aerobic and combined aerobic + resistance training yielded similar reductions in liver fat

**Table 7** continued

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Steenivasa Baba et al. (2006)	65 men and women aged 38–7 years with nonalcoholic steatohepatitis	Liver enzymes	3-month diet plus 30-min exercise, 5 days/week at 60–70 % of maximal heart rate	Compliant patients also showed a decrease of BMI. No control group	AST and ALT normalized only in patients complying with exercise programme
St George et al. (2009)	141 cases of steatosis aged 48–50 years, divided between moderate-intensity exercise, low-intensity exercise and control group	Liver enzymes	3-month trial, 6 vs. 3 lifestyle counselling sessions given over 10 weeks, with advice to walk 150 min/week vs. control group; 60 % of intervention groups reported increased physical activity	Cycle ergometer prediction of maximal oxygen intake. Gains of hepatic function seen independently of weight loss.	Greatest improvement of ALT, GGT, AST if walking 150 min/week and/or increased predicted maximal oxygen intake
Straznický et al. (2012)	63 obese adults average age 55 years	Liver enzymes	Dietary restriction alone vs. dietary restriction plus moderate exercise (40 min at 65 % of maximal heart rate on alternate days, with only one session per week supervised)		7.6 % loss of body mass with dietary restriction alone. ALT decreased 20 % by dieting. No added advantage if exercise added to dieting
Sullivan et al. (2012)	18 obese adults with steatosis, average age 48 years	Proton magnetic resonance spectroscopy	16-week exercise (45–55 % peak oxygen intake, 30–60 min/day, 5 times/week, $n = 12$ ) vs. control ( $n = 6$ )	No change in hepatic VLDL-TG secretion or VLDL-apoB-100 secretion	Exercise reduced liver fat 10 %, but no change in body mass or % body fat
Suzuki et al. (2005)	348 males with high ALT values	Liver enzymes	Lifestyle change	Weight loss, binary classification of increase in activity. No control group	Improvement in ALT associated with weight loss >5 % and increase of activity
Tamura et al. (2005)	14 patients with type 2 diabetes mellitus	Proton magnetic resonance spectroscopy	2-week comparison, diet or diet + exercise (walking 60–90 min/day for an increase in energy expenditure of about 0.7 MJ/day)	Diet + exercise group younger than diet alone	Hepatic lipids decreased in both groups, but no added effect of exercise
Thamer et al. (2007, 2008)	48 M, 64 F, average age 46 years	Proton magnetic resonance spectroscopy	Diet + 3 h/week aerobic exercise monitored by Polar pulse-counter, average follow-up of 264 days	Decrease of liver fat associated with improved insulin sensitivity. Subjects with high adipose and liver fat control less from lifestyle. No control group intervention. No control group	33 % decrease of liver fat, despite little change in overall body fat; response less marked in alleles of peroxisome proliferator-activated receptor-gene
Thomas et al. (2006)	Ten obese adults, age not stated	Proton magnetic resonance spectroscopy, liver enzymes	Diet + pedometer monitored recommendation of 10,000 steps/day	Changes associated with loss of body fat. No control group	40 % decrease of liver fat, improvement of AST but not ALT

Table 7 continued

Author	Sample	Liver assessment	Intervention	Measures and comments	Outcome
Thompson et al. (2010)	41 men aged 45–64 years	ILK-6, ALT	24-week exercise (30 min rising to 60 min at 50 % increasing to 70 % maximal oxygen intake 3 rising to 4 days/week) + 2 weeks detraining vs. control	No comment on nutrition.	IL-6 fell with 12-week moderate exercise; lost with 2-week detraining; ALT reduced at 24 weeks but not at 12 weeks, benefit retained with 2-week detraining
Ueno et al. (1997)	25 obese (ten served as controls)	Liver enzymes and histology	Restricted diet and walking or jogging for 3 months	No control group	Improvement in AST, ALT and steatosis
van der Heijden et al. (2010b)	15 obese and 14 lean adolescents	Proton magnetic resonance spectroscopy	12-week aerobic programme (30 min sessions, 2 or 4 times/week at 70 % of maximal oxygen intake)	No control group. Exercise increased hepatic and peripheral insulin sensitivity	Hepatic fat reduced in obese but not in lean subjects, without weight loss
van Der Heijden et al. (2010a)	Obese adolescents (6 M, 6 F)	Proton magnetic resonance spectroscopy	12-week resistance exercise (twice/week, all major muscle groups)	No control group. Body mass increased 2.6 kg	Hepatic fat content unchanged, but insulin sensitivity increased 24 %

hepatic function by modulating myostatin output. Myostatin inhibits muscle growth, thus predisposing to obesity, hepatic insulin resistance and diabetes; it may also have more direct effects upon hepatocytes (Allen et al. 2011). Inactivation of the myostatin gene in mice caused hepatic steatosis in the absence of any change in muscle mass (Mukherjee et al. 2007), and injection of recombinant myostatin slowed overall growth through a decrease in IGF-1 induced AKT phosphorylation, again without change of muscle mass (Hittel et al. 2010). Finally, both mouse and human liver cell cultures developed apoptosis when incubated with recombinant activin, which binds to the same receptors as myostatin (Woodruff et al. 1993; Chen et al. 2000).

#### Exercise dose recommendations in hepatic disease

From the investigations discussed above, we may conclude that regular aerobic exercise can reduce liver fat levels and this benefit can occur, albeit probably to a lesser extent, without weight loss. In humans, the majority of therapeutic programmes have prescribed exercise at moderate to vigorous intensities for 3–5 days per week (Table 7). However, clearer information is needed on the efficacy of resistance versus aerobic exercise, the minimum dose of physical activity required for benefit, the exercise tolerance of individuals with NAFLD, and doses of exercise that may lead to hepatic injury.

Whilst it appears that regular aerobic exercise of moderate or vigorous intensity is effective in decreasing hepatic fat content, vigorous exercise may not always be practical. Fatigue (probably centrally mediated) is a frequent concomitant of hepatic steatosis (Bergasa et al. 2004), and this may reduce a patient's motivation, or even preclude participation in sustained aerobic activity, especially if this is vigorous. Moreover, co-morbid obesity can in itself reduce functional capacity and discourage involvement in exercise programmes, especially if vigorous activity is required. In this context, the only study to date that has examined predictors of physical activity adoption and adherence in a NAFLD cohort concluded that initial confidence in the ability to exercise was often low, in part because of a fear of falling (Frith et al. 2010). Whilst participation in a supervised exercise programme with individuals similar to oneself is well known to improve self-efficacy and reduce fears of falling, in patients where such an approach is found to be ineffective, more unconventional tactics may be needed to increase daily energy expenditures. One investigation demonstrated that a significant reduction of ALT could be achieved by regular voluntary and electrical stimulation of the quadriceps and hamstring muscles in individuals who were resistant to lifestyle intervention (Kawaguchi et al. 2011).

**Table 8** Animal studies of metabolic syndrome and hepatic fat content with exercise interventions

Author	Animals tested	Intervention	Comments	Outcome
Botezelli et al. (2010)	Male Wistar rats initially aged 28 weeks fed on 60 % fructose diet	Swimming 1 h/day at anaerobic threshold, starting at 28 or 90 days	Swimming improved insulin sensitivity	Liver lipids and AST corrected by early or late training, but ALT unchanged
Cameron et al. (2012)	Male Wistar rats fed high-fat/high-carbohydrate diet or corn starch	8 weeks of treadmill running, 20 min/day rising to 30 min/day, 5 days/week; 0 % incline, speed 1 km/h	Exercise decreased body fat, abdominal fat and blood glucose, improved blood lipid profile	Liver mass and hepatic fat decreased in experimental group
Charbonneau et al. (2005)	Female Sprague–Dawley rats initially 10 weeks old, fed high-fat diet	Treadmill exercise progressing to 60 min/day at 26 m/min, 10 % slope	Glucagon resistance of obese rats also prevented by exercise	28 % gain of hepatic fat with high-fat diet was completely reversed by exercise
Cintra et al. (2012)	Obese mice fed high-fat diet	8 weeks of running (50 min/day 5 days/week at 1 km/h)	Associated reduction of regulatory element-binding protein-1c	1.7-fold reduction of liver fat
Colombo et al. (2005)	Male diabetic fatty Zucker rats	5 weeks of treadmill running (1 h/day, 6 days/week, 20 m/min)	Modification of many hepatic genes associated with lipogenesis and detoxification	Liver fat decreased
Corriveau et al. (2008)	Three groups of ovariectomized vs. one group sham-operated rats	25 % dietary restriction vs. dietary restriction + resistance exercise (weighted stair climbing 5 times/week)	Abdominal fat also reduced by resistance exercise	Liver lipid accumulation not stopped by dietary restriction, but reversed by resistance exercise
Gauthier et al. (2003)	Female Sprague–Dawley rats initially 6 weeks old, fed high-fat diet	Treadmill exercise progressing to 60 min/day at 26 m/min, 10 % slope	Treadmill concentrations also reduced by exercise	Liver triacylglycerol content substantially reduced by concomitant exercise relative to sedentary controls
Gauthier et al. (2004); Yasari et al. (2009)	Female Sprague–Dawley rats initially 6 weeks old fed high-fat diet	Treadmill exercise progressing to 60 min/day at 26 m/min, 10 % slope introduced from 8th to 16th week of high-fat diet	Leptin concentrations also reduced by exercise	Exercise decreased visceral fat 30 %
Hao et al. (2010)	12-week ovariectomized female rats	Treadmill running (10–18 min/day at 0 % grade for 15–60 days)	Associated increase in HDL/total cholesterol ratio	Exercise reduced hepatic fat relative to controls
Haram et al. (2009)	Rats bred for low intrinsic running capacity	60 min/day exercise for 8 weeks, either high-intensity interval work (4 min bouts at 85–90 % maximal oxygen intake) or running same distance at continuous 70 % maximal oxygen intake	Both programmes equally effective in reducing body mass and fat mass, and in increasing insulin receptor phosphorylation in the liver	Interval training more effective in reversing metabolic syndrome
Leite et al. (2009)	Adult F Wistar rats, half of sample ovariectomized	Sedentary vs. resistance training (weighted ladder climbing), 4–9 climbs every 3 days	Other fat depots also reduced	Exercise reduced liver fat content, less in ovariectomized than in intact animals
Marques et al. (2010)	C57/BL6 mice fed standard chow or very high-fat diet	Sedentary vs. 8 weeks treadmill running (60 min/day, 5 days/week at 1 km/h)	Exercise reduced insulin resistance, cholesterol and triglycerides	Video-microscopy estimate of hepatic fat content, AST and ALT reduced in exercised animals
Moura et al. (2011)	60-day alloxan-treated diabetic Wistar rats	Swimming 1 h/day, 5 days/week for 44 days with load 3.5 % body mass (below anaerobic threshold)		Hepatic fat content lower in swimmers than in sedentary peers
Pighon et al. (2009a, b)	Ovariectomized rats following 8 weeks of food restriction + resistance exercise	Normal feeding vs. food restriction vs. resistance training (weighted ladder climbing)		Dietary restriction prevents regain of liver fat; resistance exercise also avoids regain

Table 8 continued

Author	Animals tested	Intervention	Comments	Outcome
Rector et al. (2011)	Hyperphagic, Otsuka Long–Evans Tokushima Fatty rats initially aged 4 weeks	36-week voluntary wheel running vs. sedentary controls	Exercise group showed greater benefits, including increased hepatic mitochondrial fatty acid oxidation, enhanced oxidative enzyme function and protein content, and suppression of hepatic lipogenic proteins	Exercise prevented hepatic fat accumulation (also prevented by dietary restriction)
Schultz et al. (2012)	Male C57BL/6 mice fed high-fat diet	Unweighted swimming progressing to 60 min/day		15 % reduction of steatosis relative to high-fat controls
Sene-Florese et al. (2009)	Male Wistar rats aged 90–120 days fed high-fat diet	90 min/day swimming vs. 330-min/day swimming sessions		Intermittent exercise more effective than continuous in preventing hepatic fat accumulation
Takeshita et al. (2012)	Mice initially aged 6 weeks	Access to running wheel (covering 4.9 km/day) vs. sedentary control	Associated lower plasma leptin and insulin-like growth factor-1 levels	Exercisers had lower liver weights and liver triglyceride content
Thyfault et al. (2009)	Rats selected for high (1,514 m) and low (200 m) running capacity		Histology and biochemistry examined	Unfit rats had reduced mitochondrial content, reduced oxidative capacity, increased peroxisomal activity, steatosis and fibrosis and apoptosis
Yasari et al. (2010)	Female rats given high-fat diet from 6 to 8th week	Treadmill running progressing to 60 min/day at 26 m/min, 10 % slope	Decrease in hepatic SCD-1 mRNA levels and protein content	Liver triglyceride content not affected by exercise
Yasari et al. (2006)	Female Sprague–Dawley rats fed high-fat diet	Treadmill running progressing to 60 min/day at 26 m/min, 10 % slope		If training stopped, trained protected rats for 2 weeks, but no protection 6 weeks post-exercise



**Table 9** Cellular and molecular markers of hepatic dysfunction in steatosis and insulin resistance and effects of exercise training

Manifestation	Exercise	Exercise training response	Comment	Author
	2 h of treadmill running (26 m/min, 1.5 % grade)	Increased expression of insulin-like growth factor binding protein-1 mRNA	Acute bout of exercise	Anthony et al. (2001)
	7 days of voluntary wheel running	Increased release of hepatic insulin-sensitizing substance (HISS), effective in countering age-related decrease of HISS	Leads to increased peripheral uptake of glucose	Chowdhury et al. (2011, 2013)
Activation of c-jun N-terminal kinase (JNK) and nuclear factor kappa-B (NF-κB)	8 weeks, 1 h/day, with a tail weight 5 % of body mass	Reduced phosphorylation of the two major metabolic markers of reticular stress (protein-kinase like endoplasmic reticular kinase, PERK and eukaryotic initiation factor 2, eIF2 phosphorylation)	Proinflammatory molecules	da Luz et al. (2011)
Increased activity of the hepatic ketone synthesis pathway	10 weeks of treadmill running, progressing to two 60-min bouts/day at 31 m/min	Ketone synthesis normalized, decrease in HMG-CoA (rate-limiting enzyme for cholesterol synthesis)		El Midaoui et al. (2006)
Depressed activity of phosphatidylinositol 3-kinase			Central mediating role in action of insulin on hepatocytes	Katsanos (2004)
Reduction of hepatic IGF-1	8 weeks of treadmill running progressing to 60 min/day at a speed of 31 m/min and a slope of 8 %	Reduction of hepatic IGF-1 attenuated		Leme et al. (2009)
Increased activity of alpha-ketoacid dehydrogenase (BCKDH) complex	4 weeks of treadmill running, progressing to 45 min/day at 25 m/min, 6 % slope	Activity of alpha-ketoacid dehydrogenase (BCKDH) complex normalized	Rate-limiting enzyme in the catabolism of branched-chain amino acids	Li et al. (2001)
	8 weeks of swimming, 60 min/day, 5 days/week (mouse study)	Increased hepatic expression of endosomal adaptor protein APPL1, decreased expression of TRB3, increasing hepatic response to insulin	APPL1 blocks the association of protein kinase AKT with its endogenous inhibitor tribbles-related protein 3 (TRB3)	Marinho et al. (2012)
	70–80 days of voluntary wheel running	Glucose clearance of livers less than half of sedentary controls, whether in fed or fasting state	Hindlimb glucose clearance increased by training	Mondon et al. (1980)
Increased levels of alanine amino-transferase (ALT) and especially γ-glutamyl transferase, (GGT) in humans			Increased levels of hepatic enzymes associated with poor glucose tolerance; increased GGT facilitates intracellular transport of glutathione, increasing oxidative stress	Nannipieri et al. (2005)
Hyperinsulinaemia and hyperleptinaemia	30-min uphill treadmill running at 20 m/min 4 times/week	Up-regulation of hepatic insulin-receptor substrate-2 protein		Park et al. (2007)
	36 days of voluntary wheel running (6.4–9.7 km/day)	Down-regulation of hepatic glucose kinase	Possibly caused by hypoinsulinaemia	Zawalich et al. (1982)

All studies in rats unless otherwise indicated

## Conclusions

Like many body systems, the liver seems well adapted to meet the demands of regular moderate physical activity. However, function becomes impaired with prolonged periods of inadequate physical activity, and extremely prolonged vigorous exercise can also have adverse consequences, particularly under harsh environmental conditions.

Acute exercise stimulates hepatic glycogenolysis and gluconeogenesis, increases the synthesis of some proteins, and may cause oxidative stress. Enzymes involved in carbohydrate metabolism are up-regulated, and lipogenic enzymes are down-regulated. Humoral changes seem the primary triggers for these changes, but the possible roles of hepatic afferent nerves, cytokines, reactive oxygen species, and reduced hepatic blood flow remain to be clarified.

Regular moderate exercise appears to build upon the changes induced by a single session of vigorous physical activity, although further studies are needed in individuals who began training with a low hepatic fat content. In obese subjects, hepatic fat content is reduced, hypertrophy of hepatic tissue is stimulated, and clearance functions are enhanced by exercise training. Blood glucose homeostasis is also improved because of increased glycogen storage and an up-regulation of enzymes involved in carbohydrate metabolism. Fat storage is decreased by a down-regulation of lipogenic enzymes and increased lipid metabolism. Production of heat shock proteins is increased and the secretion of orixogenic proteins is decreased. Increases of antioxidant enzymes and stores of reduced glutathione enhance resistance to oxidant stress. Triggers of metabolic responses to chronic exercise seem modulations of insulin, insulin-like growth factor, glucagon and interleukin-6.

Inadequate physical activity predisposes to steatosis and associated disorders, including the metabolic syndrome, cardiovascular disease and diabetes mellitus. Simple steatosis can progress to hepatitis, cirrhosis and even hepatic carcinoma. Therapeutic exercise programmes restore insulin sensitivity, counteract diabetes and steatosis, and may facilitate recovery from hepatitis. However, the optimal exercise prescription remains to be defined in terms of efficacy and patient acceptance.

In summary, regular moderate physical activity makes an important contribution to the maintenance of optimal liver function, and this seems one more good reason to commend daily exercise as an important part of a healthy lifestyle.

## References

- Ahlborg G, Felig P, Hagenfel L, Hendler R, Wahren J (1974) Substrate turnover during prolonged exercise in man. Splanchnic and leg metabolism of glucose, free fatty-acids, and amino-acids. *J Clin Invest* 53(4):1080–1090
- Albu JB, Hellbronn LK, Kelley DE, Smith SR, Azuma K, Berk ES, Pi-Sunyer FX, Ravussin E, Look Ahead Research Group (2010) Metabolic changes following a 1-year diet and exercise intervention in patients with type 2 diabetes. *Diabetes* 59:627–633
- Alisi A, Locatelli M, Nobili V (2010) Nonalcoholic fatty liver disease in children. *Curr Opin Clin Nutr Metab Care* 13(4):397–402
- Allen DL, Hittel DS, McPherron AC (2011) Expression and function of myostatin in obesity, diabetes, and exercise adaptation. *Med Sci Sports Exerc* 43(10):1828–1835
- Andersen H, Borre M, Jakobsen J, Andersen PH, Vilstrup H (1998) Decreased muscle strength in patients with alcoholic liver cirrhosis in relation to nutritional status, alcohol abstinence, liver function, and neuropathy. *Hepatology* 27(5):1200–1206
- Angulo P (2002) Non-alcoholic fatty liver disease. *New Engl J Med* 346:1221–1231
- Anthony TG, Anthony JC, Lewitt MS, Donovan SM, Layman DK (2001) Time course changes in IGFBP-1 after treadmill exercise and postexercise food intake in rats. *Am J Physiol* 280:E650–E656
- Aoi W, Naito Y, Hang LP, Uchiyama K, Akagiri S, Mizushima K, Yoshikawa T (2011) Regular exercise prevents high-sucrose diet-induced fatty liver via improvement of hepatic lipid metabolism. *Biochem Biophys Res Comm* 413(2):330–335
- Apple FS, McGue MK (1983) Serum enzyme changes during marathon training. *Am J Clin Pathol* 79(6):716–719
- Arase Y, Suzuki F, Ikeda K, Kumada H, Tsuji H, Kobayashi T (2009) Multivariate analysis of risk factors for the development of type 2 diabetes in nonalcoholic fatty liver disease. *J Gastroenterol* 44(10):1064–1070
- Askeew EW, Barakat H, Kuhl GL, Dohm GL (1975) Response of lipogenesis and fatty acid synthetase to physical training and exhaustive exercise in rats. *Lipids Health Dis* 10(8):491–496
- Atalay M, Oksala NKJ, Laaksonen DE, Khanna S, Nakao C, Lappalainen J, Roy S, Hanninen O, Sen CK (2004) Exercise training modulates heat shock protein response in diabetic rats. *J Appl Physiol* 97:805–811
- Aydin C, Ince E, Koiparan S, Cangul IT, Naziroglu M, Ak F (2005) Protective effects of long term dietary restriction on swimming exercise-induced oxidative stress in the liver, heart and kidney of rat. *Cell Biochem Funct* 25(2):129–137
- Baba CS, Alexander G, Kalyani B, Pandey R, Rastogi S, Pandey A, Gourdas C (2006) Effect of exercise and dietary modification on serum aminotransferase levels in patients with nonalcoholic steatohepatitis. *J Gastroenterol Hepatol* 21(1 Pt 1):191–198
- Bacchi N, Moghetti P (2013) Exercise for hepatic fat accumulation in type 2 diabetic subjects. *Int J Endocrinol* ID 309191:1–5
- Bacchi E, Negri C, Targher G, Faccioli N, Lanza M, Zoppini G, Zanolin E, Schena F, Bonora E, Moghetti P (2013) Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 randomized trial). *Hepatology* 58(4):1287–1295
- Bae JC, Suh S, Park SE, Rhee EJ, Park CY, Oh KW, Park SW, Kim SW, Hur KY, Kim JH, Lee M-S, Lee MK, Kim K-W, Lee W-Y (2012) Regular exercise is associated with a reduction in the risk of NAFLD and decreased liver enzymes in individuals with NAFLD independent of obesity in Korean adults. *PLoS One* 7(10):e46819
- Balasubramian K, Mawer GE, Simons PJ (1970) The influence of dose on the distribution and elimination of amylobarbitone in healthy subjects. *Br J Pharmacol* 40(3):578–579
- Baldwin KM, Reitman JS, Terjung RL, Winder WW, Holloszy JO (1973) Substrate depletion in different types of muscle and liver during prolonged running. *Am J Physiol* 225(5):1045–1050

- Banzet S, Koulmann N, Simler N, Chapot R, Serrurier B, Peinnequin A, Bigard X (2009) Control of gluconeogenic genes during intense/prolonged exercise: hormone independent effect of muscle-derived IL-6 on hepatic tissue and PEPCK mRNA. *J Appl Physiol* 107(6):1830–1839
- Barsalani R, Chapados NA, Lavoie J-M (2010) Hepatic VLDL-TG production and MTP gene expression are decreased in ovariectomized rats: effects of exercise training. *Hormone Metabol Res* 42(12):860–867
- Barsalani R, Riesco E, Lavoie JM, Dionne IJ (2013) Effect of exercise training and isoflavones on hepatic steatosis in overweight postmenopausal women. *Climacteric* 16(1):88–95
- Bay Nielsen H, Secher NH, Clemmesen O, Ott P (2005) Maintained cerebral and skeletal muscle oxygenation during maximal exercise in patients with liver cirrhosis. *J Hepatol* 43(2):266–271
- Beard ME, Hamer JW, Hamilton G, Maslowski AH (1979) Jogger's heat stroke. *NZ Med J* 89(631):159–161
- Behrens G, Matthews CE, Moore SC, Freedman ND, McGlynn KA, Everhart JE, Hollenbeck AR, Leitzmann MF (2013) The association between frequency of vigorous physical activity and hepatobiliary cancers in the NIH-AARP diet and health study. *Eur J Epidemiol* 28(1):55–66
- Bejma J, Ramires P, Ji LL (2000) Free radical generation and oxidative stress with ageing and exercise: differential effects in the myocardium and liver. *Acta Physiol Scand* 169(4):343–351
- Bellentani S, Saccoccio G, Masutti F, Croce LS, Brandi G, Sasso F, Cristanini G, Tiribelli C (2000) Prevalence of and risk factors for hepatic steatosis in Northern Italy. *Ann Intern Med* 132(2):112–117
- Berg A, Keul J (1982) Serum enzymkinetik während und nach intensiver Langzeitbelastung (Serum enzyme kinetics during and after endurance exercise). *Dtsch Z fur Sportmed* 33:12–17
- Bergasa NV, Mehlman J, Bir K (2004) Aerobic exercise: a potential therapeutic intervention for patients with liver disease. *Med Hypoth* 62(6):935–941
- Berglund ED, Lee-Young RS, Lustig DG, Lynes SE, Donahue EP, Camacho RC, Meredith ME, Magnuson MA, Charron MJ, Wasserman DH (2009) Hepatic energy state is regulated by glucagon receptor signaling in mice. *J Clin Invest* 119(8):2412–2422
- Berglund ED, Lustig DG, Baheza RA, Hasenour CM, Lee-Young RS, Donahue EP, Lynes SE, Swift LL, Charron MJ, Damon BM, Wasserman DH (2011) Hepatic glucagon action is essential for exercise-induced reversal of mouse fatty liver. *Diabetes* 60(11):2720–2729
- Bergstrom J, Hermansen L, Hultman E, Saltin B (1967) Diet, muscle glycogen and physical performance. *Acta Physiol Scand* 71(2):140–150
- Bermon S, Ferrari P, Bernard P, Altare S, Dolisi C (1999) Responses of total and free insulin-like growth factor-I and insulin-like growth factor binding protein-3 after resistance exercise and training in elderly subjects. *Acta Physiol Scand* 165(1):51–56
- Bexfield NA, Parcell AC, Nelson WB, Foote KM, Mack GW (2009) Adaptations to high-intensity intermittent exercise in rodents. *J Appl Physiol* 107(3):749–754
- Boel J, Andersen LB, Hansen SH, Dossing M (1984) Hepatic drug metabolism and physical fitness. *Clin Pharmacol Therap* 36(1):121–126
- Bonekamp S, Barone B, Clark JM, Stewarts KJ (2008) The effect of exercise training intervention on hepatic steatosis. *Hepatol* 48(S1):806A
- Børsheim E, Knardahl DS, Høstmark AT (1999) Short-term effects of exercise on plasma very low density lipoproteins (VLDL) and fatty acids. *Med Sci Sports Exerc* 31(4):522–530
- Botezelli JD, Mora RF, Dalia RA, Moura LP, Cambri LT, Ghezzi AC, Voltarelli FA, Mello MA (2010) Exercise counteracts fatty liver disease in rats fed on fructose- rich diet. *Lipids Health Dis* 9:116 (On-line publication)
- Bouchard C, Shephard RJ (1994) Physical activity, fitness and health: the model and key concepts. In: Bouchard C, Shephard RJ, Stephens T (eds) *Physical activity, fitness and health*. Human Kinetics, Champaign, pp 77–88
- Bozzetto L, Prinster A, Annuzzi G, Costagilola L, Mangione A, Vitelli A, Mazzarella R, Longobardo M, Mancini M, Vigorito C, Riccardi G, Rivellese AA (2012) Liver fat is reduced by isoenergetic MUFA diet in a controlled random study in type 2 diabetic patients. *Diabetes Care* 35(7):1429–1435
- Brinkley TE, Halverstadt A, Phares DA, Ferrell RE, Prigeon RL, Hagberg JM, Goldberg AP (2011) Hepatic lipase gene -514CT variant is associated with exercise training-induced changes in VLDL and HDL by lipoprotein. *J Appl Physiol* 111(6):1871–1876
- Browning JD, Szczepaniak LS, Dobbins R, Nuremberg P, Horton JD, Cohen JC, Grundy SM, Hobbs HH (2004) Prevalence of hepatic steatosis in an urban population in the United States: impact of ethnicity. *Hepatol* 40(6):1387–1395
- Bunch TW (1980) Blood test abnormalities in runners. *Mayo Clin Proc* 55(2):113–117
- Bürger-Mendonça M, Bielavsky M, Barbosa FCR (2008) Liver overload in Brazilian triathletes after half-ironman competition is related to muscle fatigue. *Ann Hepatol* 7(3):245–248
- Burneiko RC, Diniz YS, Galhardi CM, Rodrigues HG, Ebaid GM, Faine LA, Padovani CR, Cicogna AC, Novelli EL (2006) Interaction of hypercaloric diet and physical exercise on lipid profile, oxidative stress and antioxidant defenses. *Food Chem Toxicol* 44(7):1167–1172
- Busse M, Nordhusen D, Tegtbur U, Machold H, Panning B (2004) Sorbitol clearance during exercise as a measure of hepatic and renal blood flow. *Clin Sportsmed Int* 1:1–8
- Caldwell S, Lazo M (2009) Is exercise an effective treatment for NASH? Knowns and unknowns. *Ann Hepatol* 8(Suppl 1):S60–S66
- Cameron I, Alam MS, Wang J, Brown L (2012) Endurance exercise in a rat model of metabolic syndrome. *Can J Physiol Pharmacol* 90(11):1490–1497
- Campillo B, Chapelain C, Bonnet JC, Frisdal E, Devanlay M, Bouissou P, Fouet P, Wirquin E, Atlan G (1990a) Hormonal and metabolic changes during exercise in cirrhotic patients. *Metab Clin Exp* 39(1):18–24
- Campillo B, Fouet P, Bonnet JC, Atlan G (1990b) Submaximal oxygen consumption in liver cirrhosis. Evidence of severe functional aerobic impairment. *J Hepatol* 10(2):163–267
- Centis E, Marzocchi R, Di Domizio S, Ciaravella MF, Marchesini G (2010) The effect of lifestyle changes in non-alcoholic fatty liver disease. *Dig Dis* 28(1):267–273
- Cha Y-S, Kim H-Y, Daily J (2003) Exercise-trained but not untrained rats maintain free carnitine reserves during acute exercise. *Asia Pac J Clin Nutr* 12(1):120–126
- Chang S-P, Chen Y-H, Chang W-C, Liu I-M, Cheng JT (2006) Merit of physical exercise to reverse the higher gene expression of hepatic phosphoenolpyruvate carboxykinase in obese Zucker rats. *Life Sci* 79(3):240–246
- Chapados NA, Lavoie J-M (2010) Exercise training increases hepatic endoplasmic reticulum (er) stress protein expression in MTP-inhibited high-fat fed rats. *Cell Biochem Funct* 28(3):202–210
- Chapados NA, Seelaender M, Levy E, Lavoie J-M (2009) Effects of exercise training on hepatic microsomal triglyceride transfer protein content in rats. *Hormone Metabol Res* 41(4):287–293
- Charbonneau A, Couturier K, Gauthier M-S, Lavoie J-M (2005) Evidence of hepatic glucagon resistance associated with hepatic steatosis: reversal effect of training. *Int J Sports Med* 26(6):432–441

- Chen W, Woodruff TK, Mayo KE (2000) Activin A-induced HepG2 liver cell apoptosis: involvement of activin receptors and smad proteins. *Endocrinol* 141(3):1263–1272
- Chen YW, Nader GA, Baar KR, Federale MJ, Hoffman EP, Esser KA (2002) Response of rat muscle to acute resistance exercise defined by transcriptional and translational profiling. *J Physiol* 545(Pt 1):27–41
- Chen S-M, Liu C-Y, Li S-R, Huang H-T, Tsai C-Y, Jou H-J (2008) Effects of therapeutic lifestyle program on ultrasound-diagnosed nonalcoholic fatty liver disease. *J Chin Med Assoc* 71(11):551–558
- Cheung O, Sanyal AJ (2009) Recent advances in nonalcoholic fatty liver disease. *Curr Opin Gastroenterol* 25(3):230–237
- Chowdhury KK, Legare DJ, Lutt WW (2011) Insulin sensitization by voluntary exercise in aging rats is mediated through hepatic insulin sensitizing substance (HISS). *Exp Gerontol* 46(1):73–80
- Chowdhury KK, Legare DJ, Lutt WW (2013) Exercise enhancement of hepatic insulin-sensitizing substance-mediated glucose uptake in diet-induced prediabetic rats. *Br J Nutr* 109(5):844–852
- Christensen NJ, Galbo H (1983) Sympathetic nervous activity during exercise. *Annu Rev Physiol* 45:139–153
- Church TS, Kuk JL, Ross R, Priest EL, Biltoff E, Blair DN (2006) Association of cardiorespiratory fitness, body mass index, and waist circumference to nonalcoholic fatty liver disease. *Gastroenterol* 130(7):2023–2030
- Cintra DE, Ropelle ER, Vitto MF, Luciano TF, Souza DR, Engelmann J, Marques SO, Lira FS, de Pinho RA, Pauli JR, De Souza CT (2012) Reversion of hepatic steatosis by exercise training in obese mice: the role of sterol regulatory element-binding protein-1c. *Life Sci* 91(11–12):395–401
- Coggan AR, Swanson SC, Mendenhall LA, Habash DL, Kien CL (1995) Effect of endurance training on hepatic glycogenolysis and gluconeogenesis during prolonged exercise in men. *Am J Physiol* 268:E375–E383
- Coker RH, Krishna MG, Lacy DB, Bracy DP, Wasserman DH (1997) Role of hepatic alpha- and beta-adrenergic receptor stimulation on hepatic glucose production during heavy exercise. *Am J Physiol* 273:E831–E838
- Coker RH, Williams RH, Yeo SE, Kortebein PM, Bodenner DL, Kern PA, Evans WJ (2009) The impact of exercise training compared to caloric restriction on hepatic and peripheral insulin resistance in obesity. *J Clin Endocrinol Metab* 94(11):4258–4266
- Colombo M, Gregersen S, Kruhoeffner M, Agger A, Xiao J, Jeppesen PB, Orntoft T, Ploug T, Galbo H, Hermansen K (2005) Prevention of hyperglycemia in Zucker diabetic fatty rats by exercise training: effects on gene expression in insulin-sensitive tissues determined by high-density oligonucleotide microarray analysis. *Metabolism* 54(12):1571–1581
- Conjeevaram HS, Tiniakos DG (2011) Editorial: exercise for NAFLD: does intensity matter? *Am J Gastroenterol* 106(3):470–475
- Corriveau P, Paquette A, Brochu M, Prud'homme D, Rabasa-Lhoret R, Lavoie JM (2008) Resistance training prevents liver fat accumulation in ovariectomized rats. *Maturitas* 59(3):259–267
- Crevenna R, Schmidinger M, Keilani M, Nuhr M, Nur H, Zöch C, Zielinski C, Fialka-Moser V, Quittan M (2003) Aerobic exercise as additive palliative treatment for a patient with advanced hepatocellular cancer. *Wien Med Wochenschr* 153(9–10):237–240
- Cullinane E, Siconolfi S, Saritelli A, Thompson PD (1982) Acute decrease in serum triglycerides with exercise—is there a threshold for an exercise effect. *Metab Clin Exp* 31(8):844–847
- da Luz G, Frederico MJS, da Silva S, Vitto MF, Cesconetto PA, de Pinho RA, Pauli JR, Silva ASR, Cintra DE, Ropelle ER, De Souza CT (2011) Endurance exercise training ameliorates insulin resistance and reticulum stress in adipose and hepatic tissue in obese rats. *Eur J Appl Physiol* 111(9):2015–2023
- da Silva LA, Pinho CA, Rocha LG, Tuon T, Silveira PC, Pinho RA (2009) Effect of different models of physical exercise on oxidative stress markers in mouse liver. *Appl Physiol Nutr Metab* 34(1):60–65
- Daemen MJAP, Thijssen HHW, van Essen H, Vervoort-Peters HT, Prinzen FW, Struyker Boudier HA, Smits JF (1989) Liver blood flow measurement in the rat: the electromagnetic versus the microsphere and clearance methods. *J Pharmacol Methods* 21(4):287–297
- Daggan RN, Zafeiridis A, Dipla K, Puglia CD, Gratz I, Catalano E, Kendrick ZV (2000) The effects of chronic exercise on anaesthesia-induced hepatotoxicity. *Med Sci Sports Exerc* 32(12):2024–2028
- Davies KJ, Quintanilha AT, Brooks GA, Packer L (1982) Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Comm* 107(4):1198–1205
- de Araújo CC, Silva JD, Samary CS, Guimarães IH, Marques PS, Oliveira GP, do Carmo LGRR, Goldenberg RC, Bakker-Abreu I, Diaz BL, Rocha NN, Capelozzi VL, Pelosi P, Rocco PRM (2012) Regular and moderate exercise before experimental sepsis reduces the risk of lung and distal organ injury. *J Appl Physiol* 112(7):1206–1214
- De Paz JA, Villa JG, Lopez P, Gonzalez-Gallego J (1995) Effects of long-distance running on serum bilirubin. *Med Sci Sports Exerc* 27(12):1590–1594
- de Piano A, de Mello M, Sanches PL, da Silva P, Campos R, Carnier J, Corgosinho F, Foschini D, Masquio D, Tock L, Oyama L, do Nascimento C, Tufik S, D'Amaso AR (2012) Long-term effects of aerobic plus resistance training on the adipokines and neuropeptides in nonalcoholic fatty liver disease obese adolescents. *Eur J Gastroenterol Hepatol* 24(11):1313–1324
- DeLissio M, Goodyear LJ, Fuller S, Krawitt EL, Devlin JT (1991) Effects of treadmill exercise on fuel metabolism in hepatic cirrhosis. *J Appl Physiol* 70(1):210–216
- Della Corte C, Alisi A, Iorio R, Alterio A, Nobili V (2011) Expert opinion on current therapies for nonalcoholic fatty liver disease. *Expert Opin Pharmacother* 12(12):1901–1911
- Devries MC, Samjoo IA, Hamadeh MJ, Tarnopolsky MA (2008) Effect of endurance exercise on hepatic lipid content, enzymes, and adiposity in men and women. *Obesity (Silver Spring)* 16(10):2281–2288
- Dharancy S, Lemyze M, Boleslawski E, Neviere R, Declerck N, Canva V, Wallaert B, Mathurin P, Pruvot FR (2008) Impact of impaired aerobic capacity on liver transplant candidates. *Transplantation* 86(8):1077–1083
- Dohm GL, Kasperek GJ, Barakat HA (1985) Time course of changes in gluconeogenic enzyme activities during exercise and recovery. *Am J Physiol* 249(1 Pt 1):E6–E11
- Domingos MM, Rodrigues MFC, Stotzer US, Bertucci DR, Souza MKC, Marine DA, Gatto CV, de Araujo HSS, De Andrade Perez SE (2012) Resistance training restores the gene expression of molecules related to fat oxidation and lipogenesis in the liver of ovariectomized rats. *Eur J Appl Physiol* 112(4):1437–1444
- Donovan CM, Pagliassotti MJ (1990) Enhanced efficiency of lactate removal after endurance training. *J Appl Physiol* 68(3):1053–1058
- Donovan CM, Sumida KD (1990) Training improves glucose homeostasis in rats during exercise via glucose production. *Am J Physiol* 258(3 Pt 2):R770–R778
- Døssing M (1985) Effect of acute and chronic exercise on hepatic drug metabolism. *Clin Pharmacokinet* 10(5):426–431
- Dowman JK, Armstrong MJ, Tomlinson JW, Newsome PN (2011) Current therapeutic strategies in non-alcoholic fatty liver disease. *Diabetes Obes Metab* 13(8):692–702

- Dressendorfer RH, Wade CE, Hornick C, Timmis GC (1982) High-density lipoprotein-cholesterol in marathon runners during a 20-day road race. *JAMA* 247(12):1715–1717
- Drouin R, Robert G, Milot M, Massicotte D, Peronnet F, Lavoie C (2004) Swim training increases glucose output from liver perfused in situ with glucagon in fed and fasted rats. *Metab Clin Exp* 53(8):1027–1031
- Ducry JJ, Howald H, Zysset T, Bircher J (1979) Liver function in physically trained subjects. Galactose elimination capacity, plasma disappearance of ICG, and aminopyrine metabolism in long distance runners. *Dig Dis Sci* 24(3):192–196
- Duvnjak M, Tomasic V, Gomercic M, Smircic Duvnjak L, Barsic N, Lerotic I (2009) Therapy of nonalcoholic fatty liver disease: current status. *J Physiol Pharmacol* 60(Suppl 7):57–66
- Dyke TM, Sams RA, Hinchcliff KW (1998) Intensity-dependent effects of acute submaximal exercise on the pharmacokinetics of bromsulphalein in horses. *Am J Vet Res* 59(11):1481–1487
- Eckard C, Cole R, Lockwood J, Torres DM, Williams CD, Shaw JC, Harrison SA (2013) Prospective histopathologic evaluation of lifestyle modification in nonalcoholic fatty liver disease: a randomized trial. *Ther Adv Gastroenterol* 6(4):249–259
- Eklöv L, Moldéus P, Orrenius S (1984) Oxidation of glutathione during hydroperoxide metabolism: a study using isolated hepatocytes and the glutathione inhibitor 1,3-bis(2-chloroethyl)-1-nitrosourea. *Eur J Biochem* 138(3):459–463
- El Midaoui A, Chiasson JL, Tancrede G, Nadeau A (2006) Physical training reverses the increased activity of the hepatic ketone body synthesis pathway in chronically diabetic rats. *Am J Physiol* 290(2):E207–E212
- Eran H, Ben-Ari Z, Heled Y, Moran DS, Shani Y, Epstein Y (2004) Liver transplantation in exertional heat stroke: a medical dilemma. *Intensive Care Med* 30(7):1474–1478
- Eslami L, Merat S, Nasser-Moghaddam S (2009) Treatment of non-alcoholic fatty liver disease (NAFLD): a systematic review. *Middle East J Dig Dis* 1(2):89–99
- Fabbri A, Bianchi G, Zoli M, Marchi E, Marchesini G (1991) Effect of physical exercise on one-sample antipyrine clearance. *Ital J Gastroenterol* 23(2):74–76
- Fallon KE, Sivyer G, Sivyer K, Dare A (1999) The biochemistry of runners in a 1600 km ultramarathon. *Br J Sports Med* 33:264–269
- Fealy CE, Haus JM, Solomon TP, Pagadala M, Flask CA, McCullough AJ, Kirwan JP (2012) Short-term exercise reduces markers of hepatocyte apoptosis in nonalcoholic fatty liver disease. *J Appl Physiol* 113(1):1–6
- Febbraio MA, Ott P, Nielsen HB, Steensberg A, Keller C, Krstrup P, Secher NH, Pedersen BK (2002) Exercise induces hepatoplanchnic release of heat shock protein 72 in humans. *J Physiol* 544(Pt 3):957–962
- Febbraio MA, Ott P, Nielsen HB, Steensberg A, Keller C, Krstrup P, Secher NH, Pedersen BK (2003) Hepatoplanchnic clearance of interleukin-6 in humans during exercise. *Am J Physiol* 285(2):E397–E402
- Felig P, Wahren J (1971) Amino acid metabolism in exercising man. *J Clin Invest* 50(12):2703–2712
- Fiebig R, Griffiths MA, Gore MT, Baker DH, Oscai L, Ney DM, Ji LL (1998) Exercise training down-regulates hepatic lipogenic enzymes in meal-fed rats: fructose versus complex-carbohydrate diets. *J Nutr* 128(5):810–817
- Fiebig RG, Hollander JM, Ji LL (2001) Exercise down-regulates hepatic fatty acid synthase in streptozotocin-treated rats. *J Nutr* 131(9):2252–2259
- Fiebig RG, Hollander JM, Ney D, Boileau R, Jeffery E, Ji LL (2002) Training down-regulates fatty acid synthase and body fat in obese Zucker rats. *Med Sci Sports Exerc* 34(7):1106–1114
- Fintini D, Pietrobbattista A, Morino G, Cafiero G, Calzolari A, Turchetta A, Brufani C, Alisi A, Giordano U, Nobili V (2012) Energy expenditure and insulin sensitivity evaluation in obese children affected by hepatosteatosis. *Pediatr Obesity* 7(2):e4–e7
- Finucane FM, Sharp SJ, Purnlow LR, Horton K, Horton J, Savage DB, Brage S, Besson H, De Lucia Rolfe E, Sleigh A, Martin HJ, Aihie Sayer A, Cooper C, Ekelund U, Griffin SJ, Wareham NJ (2010) The effects of aerobic exercise on metabolic risk, insulin sensitivity and intrahepatic lipid in healthy older people from the Hertfordshire cohort study: a randomised controlled trial. *Diabetologia* 53(4):624–631
- Flamm SD, Taki J, Moore R, Lewis SF, Keech F, Masltais F, Ahmad M, Callahan R, Dragotakes S, Alpert N, Strauss HW (1990) Redistribution of regional and organ blood volume and effect on cardiac function in relation to upright exercise intensity in healthy human subjects. *Circulation* 81(5):1550–1559
- Foigt E, Ekelund L-G, Hultman E (1976) Enzyme activities in hepatic venous blood under strenuous physical exercise. *Pflügers Arch* 361(3):287–296
- Frank S, Somani SM, Konle M (1990) Effect of exercise on propranolol pharmacokinetics. *Eur J Clin Pharmacol* 39:391–394
- Franzese A, Vajro P, Argenziano A, Puzzielio A, Iannucci MP, Saviano MC, Brunetti F, Rubino A (1997) Liver involvement in obese children. Ultrasonography and liver enzyme levels at diagnosis and during follow-up in an Italian population. *Dig Dis Sci* 42(7):1428–1432
- Frenkl R, Gyore A, Szeberenyi S (1980) The effect of muscular exercise on the microsomal enzyme system of the rat liver. *Eur J Appl Physiol* 44(2):135–140
- Friedman JE (1994) Role of glucocorticoids in activation of hepatic PEPCK gene transcription during exercise. *Am J Physiol* 266(4 Pt 1):E560–E566
- Frith J, Day CP, Robinson L, Elliott C, Jones DEJ, Newton JL (2010) Potential strategies to improve uptake of exercise interventions in non-alcoholic fatty liver disease. *J Hepatol* 52(1):112–116
- Fritsche L, Weigert C, Haring HU, Lehmann R (2008) How insulin receptor substrate proteins regulate the metabolic capacity of the liver—implications for health and disease. *Curr Med Chem* 15(13):1316–1329
- Fritsche L, Hoene M, Lehmann R, Ellingsgaard H, Hennige AM, Pohl AK, Häring HU, Schleicher ED, Weigert C (2010) IL-6 deficiency in mice neither impairs induction of metabolic genes in the liver nor affects blood glucose levels during fasting and moderately intense exercise. *Diabetologia* 53(8):1732–1744
- Froelich JE, Strauss HW, Moore RH, McKusick KA (1988) Redistribution of visceral blood volume in upright exercise in healthy volunteers. *J Nucl Med* 29(10):1714–1718
- Fuwii M, Isikawa M, Iuchi M, Tashiro S (2005) Effect of follistatin on liver regeneration and tumor growth after portal occlusion. *Hepatogastroenterol* 52(63):833–838
- García-Pagan JC, Santos C, Barbera JA, Luca A, Roca J, Rodriguez-Roisin R, Bosch J, Rodes J (1996) Physical exercise increases portal pressure in patients with cirrhosis and portal hypertension. *Gastroenterol* 111(5):1300–1306
- Gauthier M-S, Couturier K, Latour J-G, Lavoie J-M (2003) Concurrent exercise prevents high-fat-diet-induced macrovesicular hepatic steatosis. *J Appl Physiol* 94(6):2127–2134
- Gauthier M-S, Couturier K, Charbonneau A, Lavoie J-M (2004) Effects of introducing physical training in the course of a 16-week high-fat diet regimen on hepatic steatosis, adipose tissue fat accumulation, and plasma lipid profile. *Int J Obes Relat Metab Disord* 28(8):1064–1071
- Gerber L, Otgonsuren M, Mishra A, Escheik C, Bircerdinc A, Stepanova M, Younossi ZM (2012) Non-alcoholic fatty liver disease (NAFLD) is associated with low level of physical

- activity: a population-based study. *Aliment Pharmacol Therap* 36(8):772–781
- Ghanbari-Niaki A, Khabazian BM, Hossaini-Kakhak SA, Rahbarizadeh F, Hedayati M (2007) Treadmill exercise enhances ABCA1 expression in rat liver. *Biochem Biophys Res Comm* 361(4):841–846
- Ghanbari-Niaki A, Fathi R, Kakhak SAH, Farshidi Z, Barmaki S, Rahbarizadeh F, Kraemer RR (2009) Treadmill exercise's reduction of Agouti-related protein expression in rat liver. *Int J Sport Nutr Ex Metab* 19(5):473–484
- Giercksky T, Boberg KM, Farstad IN, Halvorsen S, Schrumph E (1999) Severe liver failure in exertional heat stroke. *Scand J Gastroenterol* 34(8):8245–8827
- Godin DV, Garnett ME (1992) Species related variations in anti-oxidant status- II. Differences in susceptibility to oxidative challenge. *Comp Biochem Physiol* 103(3):743–748
- Gonzalez B, Manso R (2004) Induction, modification and accumulation of HSP70s in the rat liver after acute exercise: early and late responses. *J Physiol* 556(Pt 2):369–385
- Goodpaster BH, Delany JP, Otto AD, Kuller L, Vockley J, South-Paul JE, Thomas SB, Brown J, McTigue K, Hames KC, Lang W, Jakicic JM (2010) Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults: a randomized trial. *JAMA* 304(16):1795–1802
- Gore M, Fiebig R, Hollander J, Leeuwenburgh C, Ohno H, Ji LL (1998) Endurance training alters antioxidant enzyme gene expression in rat skeletal muscle. *Can J Physiol Pharmacol* 76(12):1139–1145
- Gorski J, Oscai LB, Palmer WK (1990) Hepatic lipid metabolism in exercise and training. *Med Sci Sports Exerc* 22(2):213–221
- Griffiths MA, Fiebig RG, Gore MT, Baker DH, Esser K, Oscai L, Ji LL (1996) Exercise down-regulates hepatic lipogenic enzymes in food-deprived and re-fed rats. *J Nutr* 126(8):1959–1971
- Grønbaek H, Lange A, Birkebæk NH, Holland-Fischer P, Solvig J, Hørlyck A, Kristensen K, Rittig S, Vilstrup H (2012) Effect of a 10-week weight loss camp on fatty liver disease and insulin sensitivity in obese Danish children. *J Pediatr Gastroenterol Nutr* 54(2):223–228
- Gündüz F, Sentürk UK, Kuru O, Aktekin B, Aktekin MR (2004) The effect of one year's swimming exercise on oxidant and antioxidant capacity in aged rats. *Physiol Res* 53(2):171–176
- Haase TN, Ringholm S, Leick L, Bienso RS, Kiilerich K, Johansen S, Nielsen MM, Wojtaszewski JF, Hidalgo J, Pedersen PA, Pilegaard H (2011) Role of PGC-1 in exercise and fasting-induced adaptations in mouse liver. *Am J Physiol* 301(5):R1501–R1509
- Hallsworth K, Fattakhova G, Hollingsworth K, Thoma C, Moore S, Taylor R, Day C, Trenell M (2011) Resistance exercise reduces liver fat and its mediators in non-alcoholic fatty liver disease independent of weight loss. *Gut* 60(9):1278–1283
- Halverstadt A, Phares DA, Wilund KR, Goldberg AP, Hagberg JM (2007) Endurance exercise training raises high-density lipoprotein cholesterol and lowers small low-density lipoprotein and very low-density lipoprotein independent of body fat phenotypes in older men and women. *Metab Clin Exp* 56(4):444–450
- Hammouda O, Chtourou H, Chaouchi A, Chahed H, Ferchichi S, Kallel C, Chamari K, Souissi N (2012) Effect of short-term maximal exercise on biochemical markers of muscle damage, total antioxidant status and homocysteine levels in football players. *Asian J Sports Med* 3(4):239–246
- Hannukainen JC, Nuutila P, Borra R, Kaprio J, Kujala UM, Janatuinen T, Heinonen OJ, Kapanen J, Viljanen T, Haaparanta M, Ronnema T, Parkkola R, Knuuti J, Kalliokoski KK (2007) Increased physical activity decreases hepatic free fatty acid uptake: a study in human monozygotic twins. *J Physiol* 578(Pt 1):347–358
- Hansen J, Brandt C, Nielsen AR, Hojman P, Whitham M, Febbraio MA, Pedersen BK, Plomgaard P (2011) Exercise induces a marked increase in plasma follistatin: evidence that follistatin is a contraction-induced hepatokine. *Endocrinol* 152(1):164–171
- Hao L, Wang Y, Duan Y, Bu S (2010) Effects of treadmill exercise training on liver fat accumulation and estrogen receptor alpha expression in intact and ovariectomized rats with or without estrogen replacement treatment. *Eur J Appl Physiol* 109(5):879–886
- Haram PM, Kemi OJ, Lee SJ, Bendheim MO, Al-Share QY, Waldum HL, Gilligan LJ, Koch LG, Britton SL, Najjar SM, Wisloff U (2009) Aerobic interval training vs. continuous moderate exercise in the metabolic syndrome of rats artificially selected for low aerobic capacity. *Cardiovasc Res* 81(4):723–732
- Hassanein T, Razack A, Gavalier JS, Van Thiel DH (1992) Heatstroke: its clinical and pathological presentation, with particular attention to the liver. *Am J Gastroenterol* 87(10):1382–1389
- Hattar LN, Wilson TA, Tabotabo LA, Smith OB, Abrams SH (2011) Physical activity and nutrition attitudes in obese Hispanic children with non-alcoholic steatohepatitis. *World J Gastroenterol* 17(39):4396–4403
- Haufe S, Engeli S, Budziarek P, Utz W, Schulz-Menger J, Hermsdorf M, Wiesner S, Otto C, Haas V, de Greiff A, Luft FC, Boschmann M, Jordan J (2010) Cardiorespiratory fitness and insulin sensitivity in overweight or obese subjects may be linked through intrahepatic lipid content. *Diabetes* 59(7):1640–1647
- Havel RJ, Kane JP, Balasse EO, Segel N, Basso LV (1970) Splanchnic metabolism of free fatty acids and production of triglycerides of very low density lipoproteins in normotriglyceridemic and hypertriglyceridemic humans. *J Clin Invest* 49(11):2017–2035
- He Y, Zhang H, Fu F (2008) The effects of swimming exercise on high-fat-diet-induced steatohepatitis. *J Sports Med Phys Fitness* 48(2):259–265
- Heinrich PC, Behrmann I, Haan S, Hermanns HM, Muller-Newen G, Schaper F (2003) Principles of interleukin (IL)-6 type cytokine signalling and its regulation. *Biochem J* 374(Pt 1):1–20
- Helge JW, Watt PW, Richter EA, Rennie MJ, Kiens B (2001) Fat utilization during exercise: adaptation to a fat-rich diet increases utilization of plasma fatty acids and very low density lipoprotein-triacylglycerol in humans. *J Physiol* 537(Pt 3):1009–1020
- Hickman IJ, Jonsson JR, Prins JB, Ash S, Purdie DM, Clouston AD, Powell EE (2004) Modest weight loss and physical activity in overweight patients with chronic liver disease results in sustained improvements in alanine aminotransferase, fasting insulin, and quality of life. *Gut* 53(3):413–419
- Hittel DS, Axelson M, Sarna N, Shearer J, Huffman KM, Kraus WE (2010) Myostatin decreases with aerobic exercise and associates with insulin resistance. *Med Sci Sports Exerc* 42(11):2023–2029
- Ho KP, Xiao DS, Ke Y, Oian ZM (2001) Exercise decreases cytosolic aconitase activity in the liver, spleen and bone marrow in rats. *Biochem Biophys Res Comm* 282(1):264–267
- Hoene M, Weigert C (2010) The stress response of the liver to physical exercise. *Ex Immunol Rev* 16:163–183
- Hoene M, Lehmann R, Hennige AM, Pohl AK, Häring HU, Schleicher ED, Weigert C (2009) Acute regulation of metabolic genes and insulin receptor substrates in the liver of mice by one single bout of treadmill exercise. *J Physiol* 587(Pt 1):241–252
- Hoene M, Franken H, Fritsche L, Lehmann R, Pohl AK, Häring H-U, Zell A, Schelischer ED, Weigert C (2010) Activation of the mitogen-activated protein kinase (MAPK) signalling pathway in the liver of mice is related to plasma glucose levels after acute exercise. *Diabetologia* 53(6):1131–1141
- Holly RG, Barnard RJ, Rosenthal M, Applegate E, Pritikin N (1986) Triathlete characterization and response to prolonged strenuous competition. *Med Sci Sports Exerc* 18(1):123–127
- Holt HN, Wild SH, Wareham N, Ekelund U, Umpleby M, Shojaae-Moradie F, Holt RIG, Phillips DI, Byrne CD (2007) Differential

- effects of fatness, fitness and physical activity energy expenditure on whole-body, liver and fat insulin sensitivity. *Diabetologia* 50(8):1698–1706
- Hong H, Johnson P (1995) Antioxidant enzyme activities and lipid peroxidation levels in exercised and hypertensive rat tissues. *Int J Biochem Cell Biol* 27(9):923–931
- Hsieh SD, Yoshinaga H, Muto T, Sakurai Y (1998) Regular physical activity and coronary risk factors in Japanese men. *Circulation* 97(7):661–665
- Hu Y, Gursoy E, Cardounel A, Kalimi M (2000) Biological effects of single and repeated swimming stress in male rats: beneficial effects of glucocorticoids. *Endocrine* 13(1):123–129
- Hu C, Hoene M, Zhao X, Häring HU, Schleicher E, Lehmann R, Han X, Xu G, Weigert C (2010) Lipidomics analysis reveals efficient storage of hepatic triacylglycerides enriched in unsaturated fatty acids after one bout of exercise in mice. *PLoS One* 5(10):e13318
- Huang H, Iida KT, Sone H, Ajisaka R (2007) The regulation of adiponectin receptor expression by acute exercise in mice. *Exp Clin Endocrinol Diabetes* 115(7):417–422
- Huang C-C, Tsai S-C, Lin W-T (2008) Potential ergogenic effects of L-arginine against oxidative and inflammatory stress induced by acute exercise in aging rats. *Exp Gerontol* 43(6):571–577
- Huang C-C, Lin W-T, Hsu F-L, Tsai PW, Hou CC (2010) Metabolomics investigation of exercise-modulated changes in metabolism in rat liver after exhaustive exercise. *Eur J Appl Physiol* 108(3):557–566
- Huang CC, Yang S-C, Chan C-C, Lin W-T (2013) Ganoderma tsugae hepatoprotection against exhaustive exercise-induced liver injury in rats. *Molecules* 18(2):1741–1754
- Hurren NM, Balanos GM, Blannin AK (2011) Is the beneficial effect of prior exercise on postprandial lipaemia partly due to redistribution of blood flow? *Clin Sci* 120(12):537–548
- Jackson KC, Wohlers LM, Valencia AP, Cilenti M, Borengasser SJ, Thyfault JP, Spangenburg EE (2011) Wheel running prevents the accumulation of monounsaturated fatty acids in the liver of ovariectomized mice by attenuating changes in SCD-1 content. *Appl Physiol Nutr Metab* 36(6):798–810
- Jakovljevic D, Hallsworth K, Zalewski P, Thoma C, Klawe J, Day C, Newton J, Trenell M (2013) Resistance exercise improves autonomic regulation at rest and haemodynamic response to exercise in non-alcoholic fatty liver disease. *Clin Sci* 125(3):143–149
- James DE, Kraegen EW (1984) The effect of exercise training on glycogen, glycogen synthase and phosphorylase in muscle and liver. *Eur J Appl Physiol* 52(3):276–281
- Jenkins NT, Hagberg JM (2011) Aerobic training effects on glucose tolerance in prediabetic and normoglycemic humans. *Med Sci Sports Exerc* 43(12):2231–2240
- Jensen MD (1995) Gender differences in regional fatty-acid metabolism before and after meal ingestion. *J Clin Invest* 96(5):2297–2303
- Ji LL, Dillon D, Wu E (1990) Alteration of antioxidant enzymes with aging in skeletal muscle and liver. *Am J Physiol* 258(4 Pt 2):R918–R923
- Jiang Y, Cyress AM, Muse ED, Wu CR, Unson CG, Merrifield RB, Sakmar TP (2001) Glucagon receptor activates extracellular signal-regulated protein kinase 1/2 via cAMP-dependent protein kinase. *Proc Natl Acad Sci USA* 98(18):10102–10107
- Jin YJ, Kim KM, Hwang S, Lee SG, Ha TY, Song GW, Jung DH, Kim KH, Yu E, Shim JH, Lim YS, Lee HC, Chung YH, Lee YS, Suh DJ (2012) Exercise and diet modification in non-obese non-alcoholic fatty liver disease: analysis of biopsies of living liver donors. *J Gastroenterol Hepatol* 27(8):1341–1347
- Johnson NA, George J (2010) Fitness versus fatness: moving beyond weight loss in nonalcoholic fatty liver disease. *Hepatology* 52(1):370–381
- Johnson NA, Sachinwalla T, Walton DW, Smith K, Armstrong A, Thompson MW, George J (2009) Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Hepatology* 50(4):1105–1112
- Johnson NA, van Overbeek D, Chapman PG, Thompson MW, Sachinwalla T, George J (2012) Effect of prolonged exercise and pre-exercise dietary manipulation on hepatic triglycerides in trained men. *Eur J Appl Physiol* 112(5):1817–1825
- Jones JC, Coombes JS, Macdonald GA (2012) Exercise capacity and muscle strength in patients with cirrhosis. *Liver Transpl* 18(2):146–151
- Kang JH, Greenon JK, Omo JT, Chao C, Peterman D, Anderson L, Foess-Wood L, Sherbondy MA, Conjeevaram HS (2006) Metabolic syndrome is associated with greater histologic severity, higher carbohydrate, and lower fat diet in patients with NAFLD. *Am J Gastroenterol* 101(10):2247–2253
- Kantartzis K, Thamer C, Peter A, Machann J, Schick F, Schraml C, Konigsrainer A, Konigsrainer I, Krober S, Niess A, Fritsche A, Haring H-U, Stefan N (2009) High cardiorespiratory fitness is an independent predictor of the reduction in liver fat during a lifestyle intervention in non-alcoholic fatty liver disease. *Gut* 58(9):1281–1288
- Katsanos CS (2004) Lipid-induced insulin resistance in the liver: role of exercise. *Sports Med* 34(14):955–965
- Katz M, Bergman EN (1969) Simultaneous measurements of hepatic and portal venous blood flow in the sheep and dog. *Am J Physiol* 216(4):946–952
- Kawaguchi T, Shiba N, Maeda T, Matsugaki T, Takano Y, Itou M, Sakata M, Taniguchi E, Nagata K, Sata M (2011) Hybrid training of voluntary and electrical muscle contractions reduces steatosis, insulin resistance, and IL-6 levels in patients with NAFLD: a pilot study. *J Gastroenterol* 46(6):746–757
- Kawanishi N, Yano H, Mizokami T, Takahashi M, Oyanagi E, Suzuki K (2012) Exercise training attenuates hepatic inflammation, fibrosis and macrophage infiltration during diet induced-obesity in mice. *Brain Behav Immun* 26(6):931–941
- Kayashima S, Ohno H, Fujioka T, Taniguchi N, Nagata N (1995) Leucocytosis as a marker of organ damage induced by chronic strenuous physical exercise. *Eur J Appl Physiol* 70(5):413–420
- Kayatekin BM, Gonenc S, Acikgoz O, Uysal N, Dayi A (2002) Effects of sprint exercise on oxidative stress in skeletal muscle and liver. *Eur J Appl Physiol* 87(2):141–144
- Keating SE, Hackett DA, George J, Johnson NA (2012) Exercise and non-alcoholic fatty liver disease: a systematic review and meta-analysis. *J Hepatol* 57(1):157–166
- Kelley GA, Kelley KS, Tran ZV (2005) Aerobic exercise, lipids and lipoproteins in overweight and obese adults: a meta-analysis of randomized controlled trials. *Int J Obesity* 29(8):881–893
- Kelley GA, Kelley KS, Franklin B (2006) Aerobic exercise and lipids and lipoproteins in patients with cardiovascular disease: a meta-analysis of randomized controlled trials. *J Cardiopulm Rehabil* 26(3):131–139
- Kelly M, Keller C, Avilucea PR, Keller P, Luo Z, Xiang X, Giralt M, Hidalgo J, Saha AK, Pedersen BK, Ruderman NB (2004) AMPK activity is diminished in tissues of IL-6 knockout mice: the effect of exercise. *Biochem Biophys Res Comm* 320(2):449–454
- Kemme MJ, Burggraaf J, Schoemaker RC, Paulson S, Karim A, Lentjes EG, Childs A, Braekman RA, Cohen AF (2000) The influence of reduced liver blood flow on the pharmacokinetics and pharmacodynamics of recombinant tissue factor pathway inhibitor. *Clin Pharmacol Ther* 67(5):504–511
- Khanna S, Atalay M, Lodge JK, Laaksonen DE, Roy S, Hanninen O, Packer L, Sen CK (1998) Skeletal muscle and liver lipoyllysine content in response to exercise, training and dietary alpha-lipoic acid supplementation. *Biochem Mol Biol Intern* 46(2):297–306

- Khazaenia T, Ramsey AA, Tam YK (2000) The effects of exercise on the pharmacokinetics of drugs. *J Pharm Pharm Sci* 3(3):292–302
- Kiens B (1993) Skeletal muscle substrate utilization during submaximal exercise in man: effect of endurance training. *J Physiol* 469:459–478
- Kindermann W, Salas-Fraire, Sroka G, Miller U (1983) Serum enzymverhalten nach körperlicher Belastung—Abgrenzung von krankheitsbedingten Veränderungen (serum enzyme content after physical activity- demarcation of disease-related changes). *Herz* 3:117–123
- Kiraly MA, Campbell J, Park EB, Bates HE, Yue JTY, Rao VM, Matthews SG, Bikopoulos G, Rozakis-Adcock M, Giacca A, Vranic M, Riddell MC (2010) Exercise maintains euglycemia in association with decreased activation of c-Jun NH2-terminal kinase and serine phosphorylation of IRS-1 in the liver of ZDF rats. *Am J Physiol* 298(3):E671–E682
- Kistler KD, Brunt EM, Clark JM, Diehl AM, Sallis JF (2011) Physical activity recommendations, exercise intensity, and histological severity of nonalcoholic fatty liver disease. *Am J Gastroenterol* 106(3):470–475
- Kjaer M (1998) Hepatic glucose production during exercise. *Adv Exp Med Biol* 441:117–127
- Kjaer M, Engfred K, Fernandes A, Secher NH, Galbo H (1993) Regulation of hepatic glucose production during exercise in humans: role of sympathoadrenergic activity. *Am J Physiol* 265(2 Pt 1):E275–E283
- Klein S, Peters EJ, Holland OB, Wolfe RR (1989) Effect of short-term and long-term beta-adrenergic-blockade on lipolysis during fasting in humans. *Am J Physiol* 257(1):E65–E73
- Klotz U, Lücke C (1978) Physical exercise and disposition of diazepam. *Br J Clin Pharmacol* 5(4):349–350
- Koot BGP, van der Baan-Slootweg OH, Tamminga-Smeulders CLJ, Rijcken THP, Pels TH, Korevaar JC, van Aalderen WM, Jansen PLM, Benninga MA (2011) Lifestyle intervention for non-alcoholic fatty liver disease: prospective cohort study of its efficacy and factors related to improvement. *Arch Dis Child* 96(7):669–674
- Korenblat KM, Fabbri E, Mohammed BS, Klein S (2008) Liver, muscle, and adipose tissue insulin action is directly related to intrahepatic triglyceride content in obese subjects. *Gastroenterol* 134(5):1369–1375
- Kotroń A, Juurinen L, Tiikkainen M, Vehkavaara S, Yki-Järvinen H (2008) Increased liver fat, impaired insulin clearance, and hepatic and adipose tissue insulin resistance in type 2 diabetes. *Gastroenterol* 135(1):122–130
- Koutedakis Y, Raafat A, Sharp NC, Rosmarin MN, Beard MJ, Robbins SW (1993) Serum enzyme activities in individuals with different levels of physical fitness. *J Sports Med Phys Fitness* 33(3):252–257
- Koyama K, Kaya M, Ishigaki T, Tsujita J, Hori S, Seino T, Kasugai A (1999) Role of xanthine oxidase in delayed lipid peroxidation in rat liver induced by acute exhausting exercise. *Eur J Appl Physiol* 80(1):28–33
- Krasnoff JB, Painter PL, Wallace JP, Bass NM, Merriman RB (2008) Health-related fitness and physical activity in patients with non-alcoholic fatty liver disease. *Hepatology* 47(4):1158–1166
- Kratz A, Lewandrowski KB, Siegel AJ, Chun KY, Flood JG, Van Cott EM, Lee-Lewandrowski E (2002) Effect of marathon running on hematologic and biochemical laboratory parameters, including cardiac markers. *Am J Clin Pathol* 118(6):856–863
- Kraus WE, Houmard JA, Duscha BD, Knetzer KJ, Wharton MB, McCartney JS, Bales CW, Henes S, Samsa GP, Otvos JD, Kulkarni KR, Slentz CA (2002) Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 347(19):1483–1492
- Kregel KC, Moseley PL (1996) Differential effects of exercise and heat stress on liver HSP70 accumulation with aging. *J Appl Physiol* 80(2):547–551
- Kugelmass M, Hill DB, Vivian B, Marsano L, McCalin CJ (2003) Cytokines and NASH: a pilot study of the effects of lifestyle modification and vitamin E. *Hepatology* 38(2):413–419
- Kuk JL, Nichaman MZ, Church TS, Blair DN, Ross R (2004) Liver fat is not a marker of metabolic risk in lean premenopausal women. *Metabolism* 53(8):1066–1071
- Larson-Meyer DE, Newcomer BR, Heilbrunn LK, Volaufova J, Smith SR, Alfonso AJ, Lefevre M, Rood JC, Williamson DA, Ravussin E, Pennington CALERIE Team (2008) Effect of 6-month calorie restriction and exercise on serum and liver lipids and markers of liver function. *Obesity (Silver Spring)* 16(6):1355–1362
- Latour MG, Brault A, Huet PM, Lavoie JM (1999) Effects of acute physical exercise on hepatocyte volume and function in rat. *Am J Physiol* 276(5 Pt 2):R1258–R1264
- Lavoie JM (2002) The contribution of afferent signals from the liver to metabolic regulation during exercise. *Can J Physiol Pharmacol* 80(11):1035–1044
- Lavoie JM, Gauthier MS (2006) Regulation of fat metabolism in the liver: link to non-alcoholic hepatic steatosis and impact of physical exercise. *Cell Mol Life Sci* 63(12):1393–1409
- Lavoie JM, Fillon Y, Couturier K, Corriveau P (2002) Evidence that the decrease in liver glycogen is associated with the exercise-induced decrease in IGFBP-1. *J Appl Physiol* 93(2):798–804
- Lawlor DA, Sattar N, Smith GD, Ebrahim S (2005) The associations of physical activity and adiposity with alanine aminotransferase and gamma glutamyltransferase. *Am J Epidemiol* 161(11):1081–1088
- Lazo M, Solga SF, Horska A, Bonekamp S, Diehl AM, Brancati FL, Clark JM, Look Ahead Research Group (2008) The effect of one year of intensive lifestyle intervention on hepatic steatosis. *Hepatology* 48:813A
- Lee KY, Kim SJ, Cha YS, So JR, Park JS, Kang KS, Chon TW (2006) Effect of exercise on hepatic gene expression in an obese mouse model using cDNA microarrays. *Obesity (Silver Spring)* 14(8):1294–1302
- Lee YS, Kek BLK, Poh LKS, Saw SM, Loke KY (2008) Association of raised liver transaminases with physical inactivity, increased waist-hip ratio, and other metabolic morbidities in severely obese children. *J Pediatr Gastroenterol Nutr* 47(2):172–178
- Lee S, Bacha F, Hannon T, Kuk JL, Boesch C, Arslanian S (2012) Effects of aerobic versus resistance exercise without caloric restriction on abdominal fat, intrahepatic lipid, and insulin sensitivity in obese adolescent boys: a randomized, controlled trial. *Diabetes* 61(11):2787–2795
- Légaré A, Drouin R, Milot M, Massicotte D, Peronnet F, Massicotte G, Lavoie C (2001) Increased density of glucagon receptors in liver from endurance-trained rats. *Am J Physiol* 280(1):E193–E196
- Leite RD, Prestes J, Bernardes CF, Shigemoto GE, Pereira GB, Duarte JO, Domingos MM, Baldissera V, de Perez ASE (2009) Effects of ovariectomy and resistance training on lipid content in skeletal muscle, liver, and heart; fat depots; and lipid profile. *App Physiol Nutr Metab* 34(6):1079–1086
- Leme JACA, Silveira RF, Gomes RJ, Moura RF, Sibuya CA, Mello MAR, Luciano E (2009) Long-term physical training increases liver IGF-I in diabetic rats. *Growth Horm IGF Res* 19(3):262–266
- Lemon PW, Nagle FJ (1981) Effect of exercise on protein and amino acid metabolism. *Med Sci Sports Exerc* 13(3):141–149
- Leskinen T, Sipilä S, Alen M, Cheng S, Pietiläinen KH, Usenius JP, Suominen H, Kovanen V, Kainukainen H, Kaprio J, Kujala UM (2009) Leisure-time physical activity and high-risk fat: a



- longitudinal population-based twin study. *Int J Obes (Lond)* 33(11):1211–1218
- Leu JI, George DL (2007) Hepatic IGFBP-1 is a prosurvival factor that binds BAK, protects the liver from apoptosis, and antagonizes the proapoptotic actions of p53 at mitochondria. *Genes Dev* 21(23):3095–3109
- Levinger I, Goodman C, Peake J, Garnham A, Hare DL, Jerums G, Selig S (2009) Inflammation, hepatic enzymes and resistance training in individuals with metabolic risk factors. *Diabetes Med* 26(3):220–227
- Li Z, Murakami T, Nakai N, Nagasaki M, Obayashi M, Xu M, Sato J, Oshida Y, Sato Y, Shimomura Y (2001) Modification by exercise training of activity and enzyme expression of hepatic branched-chain alpha-ketoacid dehydrogenase complex in streptozotocin-induced diabetic rats. *J Nutr Sci Vitaminiol (Tokyo)* 47(5):345–350
- Linden MA, Meers GM, Ruebl ML, Jenkins NT, Booth FW, Laughlin MH, Ibdah JA, Thyfault JP, Reiter RS (2013) Hepatic steatosis development with 4 weeks of physical inactivity in previously active, hyperphagic OLETF rats. *Am J Physiol Regul Integr Comp Physiol* 304(9):R763–R771
- Lindfeldt J, Balkan B, van Dijk G, Scheurink A, Ahren B, Steffens AB (1993) Influence of peri-arterial hepatic denervation on the glycemic response to exercise in rats. *J Auton Nerv Syst* 44(1):45–52
- Lippi G, Schena F, Montagnana M, Salvagno GL, Banfi G, Guidi GC (2011) Significant variation of traditional markers of liver injury after a half-marathon run. *Eur J Int Med* 22(5):e36–e38
- Litvinova L, Viru A (1995) Effect of exercise and adrenal insufficiency on urea production in rats. *Eur J Appl Physiol* 70(6):536–540
- Liu J, Yeo HC, Overvik-Douki E, Hagen T, Doniger SJ, Chyu DW, Brooks GA, Ames BN (2000) Chronically and acutely exercised rats: biomarkers of oxidative stress and endogenous anti-oxidants. *J Appl Physiol* 89(1):21–28
- Long D, Blake M, McNaughton L, Angle B (1990) Hematological and biochemical changes during a short triathlon competition in novice triathletes. *Eu J Appl Physiol* 61(1–2):93–99
- Loniewski I, Sawrymowicz M, Pawlik A, Wojcicki J, Drozdziak M (2001) Lack of effect of physical exercise on pharmacokinetics of acetaminophen tablets in healthy subjects. *Acta Pol Pharm* 58(2):141–144
- Lundbergh P, Strandell T (1974) Changes in hepatic circulation at rest, during and after exercise in young males with infectious hepatitis compared with controls. *Acta Med Scand* 196(4):315–325
- Mager DR, Patterson C, So S, Rogenstein CD, Wykes LJ, Roberts EA (2010) Dietary and physical activity patterns in children with fatty liver. *Eur J Clin Nutr* 64(6):628–635
- Magkos F, Patterson BW, Mohammed BS, Mittendorfer B (2007) A single 1-h bout of evening exercise increases basal FFA flux without affecting VLDL-triglyceride and VLDL-apolipoprotein B-100 kinetics in untrained lean men. *Am J Physiol* 292(6):E1568–E1574
- Magkos F, Patterson BW, Mohammed BS, Mittendorfer B (2009) Basal adipose tissue and hepatic lipid kinetics are not affected by a single exercise bout of moderate duration and intensity in sedentary women. *Clin Sci (Lond)* 116(4):327–334
- Malhi N, Kaufman RJ (2011) Endoplasmic reticulum stress in liver disease. *J Hepatol* 54(4):795–809
- Malinoski FJ (1992) Strenuous exercise simulating hepatic injury during vaccine trials. *Vaccine* 10(1):39–42
- Margaritis I, Tessier F, Richard MJ, Marconnet P (1997) No evidence of oxidative stress after a triathlon race in highly trained competitors. *Int J Sports Med* 18(3):186–190
- Marinho R, Ropelle ER, Cintra DE, De Souza CT, Da Silva ASR, Bertoli FC, Colantonio E, D’Almeida V, Pauli JR (2012) Endurance exercise training increases APPL1 expression and improves insulin signaling in the hepatic tissue of diet-induced obese mice, independently of weight loss. *J Cell Physiol* 227(7):2917–2926
- Marques CMM, Motta VF, Torres TS, Aguila MB, Mandarim-de-Lacerda CA (2010) Beneficial effects of exercise training (treadmill) on insulin resistance and nonalcoholic fatty liver disease in high-fat fed C57BL/6 mice. *Braz J Med Biol Res* 43(5):467–475
- Martin WH, Dalsky GP, Hurley BF, Matthews DE, Bier DM, Hagberg JM, Rogers MA, King DS, Holloszy JO (1993) Effect of endurance training on plasma free fatty acid turnover and oxidation during exercise. *Am J Physiol* 265(5 Pt 1):E708–E714
- Mauriz JL, Taberero B, Garcia-Lopez J (2000) Physical exercise and improvement of liver oxidative metabolism in the elderly. *Eur J Appl Physiol* 81(1–2):62–66
- McMillan KP, Kuk JL, Church TS, Blair DN, Ross R (2007) Independent associations between liver fat, visceral adipose tissue, and metabolic risk factors in men. *Appl Physiol Nutr Metab* 32(2):265–272
- Meek SE, Nair KS, Jensen MD (1999) Insulin regulation of regional free fatty acid metabolism. *Diabetes* 48(1):10–14
- Mena P, Maynar M, Campillo JE (1996) Changes in plasma enzyme activities in professional racing cyclists. *Br J Sports Med* 30(2):122–124
- Mencin AA, Lavine JE (2011) Nonalcoholic fatty liver disease in children. *Curr Opin Clin Nutr Metab Care* 14(2):151–157
- Mikami T, Sumida S, Ishibashi Y, Ohta S (2004) Endurance exercise training inhibits activity of plasma GOT and liver caspase-3 of mice [correction of rats] exposed to stress by induction of heat shock protein 70. *J Appl Physiol* 96(5):1776–1781
- Miura H, Yoshimoto H, Kitamura S, Yamada H (2010) Fulminant hepatic failure accompanied by fatal rhabdomyolysis following exertional heatstroke. *Clin J Gastroenterol* 3:318–323
- Mondon CE, Dolkas CB, Reaven GM (1980) Site of enhanced insulin sensitivity in exercise-trained rats at rest. *Am J Physiol* 239(3):E169–E177
- Moon J, Kallman J, Winter P, Srishord M, Fang Y, Gerber L, Younossi Z (2012a) Disparities in activity level and nutrition between patients with chronic hepatitis C and blood donors. *PM R* 4(6):436–441
- Moon MK, Cho BJ, Lee YJ, Choi SH, Lim S, Park KS, Park YJ, Jang HC (2012b) The effects of chronic exercise on the inflammatory cytokines interleukin-6 and tumor necrosis factor- $\alpha$  are different with age. *Appl Physiol Nutr Metab* 37(4):631–636
- Mooy J, Arends B, Kaminade JV, Boehm R, Rahn KM, Baak MV (1986) Influence of prolonged submaximal exercise on the pharmacokinetics of verapamil in humans. *J Cardiovasc Pharmacol* 8(5):940–942
- Moses FM (1990) The effects of exercise on the gastrointestinal tract. *Sports Med* 9(3):159–172
- Moura LP, Puga GM, Beck WR, Teixeira IP, Ghezzi AC, Silva GA, Mello MAR (2011) Exercise and spirulina control non-alcoholic hepatic steatosis and lipid profile in diabetic Wistar rats. *Lipids Health Dis* 10:77 (On-line publication)
- Mukherjee A, Sidis Y, Mahan A, Raheer MJ, Xia Y, Rosen ED, Bloch KD, Thomas MK, Schneyer AL (2007) FSTL3 deletion reveals roles for TGF-beta family ligands in glucose and fat homeostasis in adults. *Proc Natl Acad Sci USA* 104(4):1348–1353
- Murakami T, Shimomura Y, Fujitsuka N, Sokabe M, Okamura K, Sakamoto S (1997) Enlargement glycogen store in rat liver and muscle by fructose-diet intake and exercise training. *J Appl Physiol* 82(3):772–775
- Musso G, Cassader M, Rosina F, Gambino R (2012) Impact of current treatments on liver disease, glucose metabolism and cardiovascular risk in non-alcoholic fatty liver disease (NAFLD):

- a systematic review and meta-analysis of randomised trials. *Diabetologia* 55(4):885–904
- Nagel D, Seiler D, Franz H, Jung K (1990) Ultra-long distance running and the liver. *Int J Sports Med* 11(6):441–445
- Nannipieri M, Gonzales C, Baldi S, Posadas R, Williams K, Haffner SM, Stern MP, Ferrannini E, Mexico Diabetes Study (2005) Liver enzymes, the metabolic syndrome, and incident diabetes the Mexico city diabetes study. *Diabetes Care* 28(7):1757–1762
- Nathwani RA, Pais S, Reynolds TB, Kaplowitz N (2005) Serum alanine aminotransferase in skeletal muscle diseases. *Hepatology* 41(2):380–382
- Navarro A, Gomez C, López-Cepero JM, Boveris A (2003) Beneficial effects of moderate exercise on mice aging: survival, behavior, oxidative stress, and mitochondrial electron transfer. *Am J Physiol* 286(3):R505–R511
- Neubauer O, König D, Kern N, Nics L, Wagner KH (2008) No indications of persistent oxidative stress in response to an ironman triathlon. *Med Sci Sports Exerc* 40(12):2119–2228
- Newton JL, Jones DEJ, Henderson E, Kane L, Wilton K, Burt AD, Day CP (2008) Fatigue in non-alcoholic fatty liver disease (NAFLD) is significant and associates with inactivity and excessive daytime sleepiness but not with liver disease severity or insulin resistance. *Gut* 57(6):807–813
- Nguyen-Duy TB, Nichaman MZ, Church TS, Blair DN, Ross R (2003) Visceral fat and liver fat are independent predictors of metabolic risk factors in men. *Am J Physiol* 284(6):E1065–E1071
- Nielsen HB, Febbraio MA, Ott P, Krstrup P, Secher NH (2007) Hepatic lactate uptake versus leg lactate output during exercise in humans. *J Appl Physiol* 103(4):1227–1233
- Nilssen O, Forde OH, Brenn T (1990) The Tromsø study. Distribution and population determinants of gamma-glutamyl transferase. *Am J Epidemiol* 132(2):318–326
- Nishida Y, Matsubara T, Tobina T, Shindo M, Tokuyama K, Tanaka K, Tanaka H (2010) Effect of low-intensity aerobic exercise on insulin-like growth factor-I and insulin-like growth factor-binding proteins in healthy men. *Int J Endocrinol* 2010.pii:452820 (On-line publication)
- Nizielski SE, Arizmendi C, Shteyngarts AR, Farrell CJ, Friedman JE (1996) Involvement of transcription factor C/EBP-beta in stimulation of PEPCK gene expression during exercise. *Am J Physiol* 270(5 Pt 2):R1005–R1012
- Noakes TD, Carter JW (1976) Biochemical parameters in athletes before and having run 160 kilometres. *S Afr Med J* 50(40):1562–1566
- Noakes TD, Carter JW (1982) The responses of plasma biochemical parameters to a 56 km race in novice and experienced ultramarathoners. *Eur J Appl Physiol* 49(2):179–186
- Nobili V, Sanyal AJ (2012) Treatment of nonalcoholic fatty liver disease in adults and children: a closer look at the arsenal. *J Gastroenterol* 47(1):29–36
- Nobili V, Manco M, Devito R, Di Commo V, Comparcola D, Sartorelli MR, Piemonte F, Marcellini M, Angulo P (2008) Lifestyle intervention and antioxidant therapy in children with nonalcoholic fatty liver disease: a randomized, controlled trial. *Hepatology* 48(1):119–128
- Nobili V, Carter-Kent C, Feldstein AE (2011) The role of lifestyle changes in the management of chronic liver disease. *BMC Medicine* 9:70 (On-line publication)
- Ochiai M, Matsuo T (2012) Increased stearoyl-CoA desaturase index and triglyceride content in the liver of rats after a single bout of swimming exercise. *Biosci Biotechnol Biochem* 76(7):1350–1355
- O'Donovan G, Thomas EL, McCarthy JP, Fitzpatrick J, Durigfhel G, Mehta S, Morin SX (2009) Fat distribution in men of different waist girth, fitness level and exercise habit. *Int J Obesity (Lond)* 33(12):1356–1362
- Ogonovszky H, Sasvari M, Dosek A, Berkes I, Kaneko T, Tahara S, Nakamoto H, Goto S, Radak Z (2005) The effects of moderate, strenuous and overtraining on oxidative stress markers and DNA repair in the rat liver. *Can J Appl Physiol* 30(2):186–195
- Ohno H, Yahata T, Yamashita K, Yamamura K, Sato Y, Yamamoto C, Hayamizu S, Taniguchi N (1988) Effect of physical training on immunoreactive gamma-glutamyltransferase in human plasma. *Enzyme* 39(2):110–114
- Orioli S, Bandinelli I, Birardi A, Chieca R, Buzzelli G, Chiarantini E (1990) Hepatic antipyrine metabolism in athletes. *J Sports Med Phys Fitness* 30(3):261–263
- Oza N, Eguchi Y, Mizuta T, Ishibashi E, Kitajima Y, Horie H, Ushirogawa M, Tsuzura T, Nakashita S, Takahashi H, Kawaguchi Y, Oda Y, Iwakiri R, Ozaki I, Eguchi T, Ono N, Fujimoto K (2009) A pilot trial of body weight reduction for nonalcoholic fatty liver disease with a home-based lifestyle modification intervention delivered in collaboration with interdisciplinary medical staff. *J Gastroenterol* 44(12):1203–1208
- Panton LB, Guillen GJ, Williams L, Graves JE, Vivas C, Cediell M, Pollock ML, Garzarella L, Krummer J, Derendorf H et al (1995) The lack of effect of aerobic exercise training on propranolol pharmacokinetics in young and elderly adults. *J Clin Pharmacol* 35(9):885–894
- Paquette AW, Wang D, Gauthier MS, Prud'homme D, Jankowski M, Gutkowska J, Lavoie JM (2007) Specific adaptations of estrogen receptor alpha and beta transcripts in liver and heart after endurance training in rats. *Mol Cell Biochem* 306(1–2):179–187
- Park S, Hong SM, Lee JE, Sung SR (2007) Chlorpromazine exacerbates hepatic insulin sensitivity via attenuating insulin and leptin signaling pathway, while exercise partially reverses the adverse effects. *Life Sci* 80(26):2428–2435
- Pedersen L, Pilegaard H, Hansen J, Brandt C, Adser H, Hidalgo J, Olesen J, Pedersen BK, Hojman P (2011) Exercise-induced liver chemokine CXCL-1 expression is linked to muscle-derived interleukin-6 expression. *J Physiol* 589(Pt 6):1409–1420
- Peeling P (2010) Exercise as a mediator of hepcidin activity in athletes. *Eur J Appl Physiol* 110(5):877–883
- Peijie C, Zicai D, Haowen X, Renbao X (2004) Effects of chronic and acute training on glucocorticoid receptors concentrations in rats. *Life Sci* 75(11):1303–1311
- Perseghin G, Lattuada G, De Cobelli F, Ragona F, Ntali G, Esposito A, Belloni E, Canu T, Terruzzi I, Scifo P, Del Maschio A, Luzi L (2007) Habitual physical activity is associated with intrahepatic fat content in humans. *Diabetes Care* 30(3):683–688
- Petersen KF, Dufour S, Befroy D, Lehrke M, Hendler RE, Shulman GI (2005) Reversal of nonalcoholic hepatic steatosis, hepatic insulin resistance, and hyperglycemia by moderate weight reduction in patients with type 2 diabetes. *Diabetes* 54(3):603–608
- Peterson KF, Price TB, Bergeron R (2004) Regulation of net hepatic glycogenolysis and gluconeogenesis during exercise: impact of type I diabetes. *J Clin Endocrinol Metab* 89(9):4656–4664
- Petridou A, Nikolaidis MG, Matsakas A, Schulz T, Michna H, Mougios V (2005) Effect of exercise training on the fatty acid composition of lipid classes in rat liver, skeletal muscle, and adipose tissue. *Eur J Appl Physiol* 94(1–2):84–92
- Pettersson J, Hindorf U, Persson P, Bengtsson T, Malmqvist U, Werkstrom V, Ekelund M (2008) Muscular exercise can cause highly pathological liver function tests in healthy men. *Br J Clin Pathol* 65(2):253–259
- Pieber K, Crevenna R, Nuhr MJ, Quittan M, Peck-Radosavljevic M, Fialka-Moser V, Wiesinger GF (2006) Aerobic capacity, muscle strength and health-related quality of life before and after

- orthotopic liver transplantation: preliminary data of an Austrian transplantation centre. *J Rehab Med* 38(5):322–328
- Pighon A, Paquette A, Barsalani R et al (2009a) Substituting food restriction by resistance training prevents liver and body fat regain in ovariectomized rats. *Climacteric* 12(2):153–164
- Pighon A, Paquette A, Barsalania R, Chapadosa NA, Rabasa-Lhoret R, Yasari S, Prud'homme D, Lavoie JM (2009b) Resistance training attenuates fat mass regain after weight loss in ovariectomized rats. *Maturitas* 64(1):52–57
- Pighon A, Gutkowska J, Jankowski M, Rabasa-Lhoret R, Lavoie JM (2011) Exercise training in ovariectomized rats stimulates estrogenic-like effects on expression of genes involved in lipid accumulation and subclinical inflammation in liver. *Metab Clin Exp* 60(5):629–639
- Pinho RA, Silva LA, Pinho CA, Scheffer DL, Souza CT, Benetti M, Carvalho T, Dal-Pizzol F (2010) Oxidative stress and inflammatory parameters after an Ironman race. *Clin J Sports Med* 20(4):306–311
- Pintus F, Mascia P (1996) Distribution and population determinants of gamma-glutamyltransferase in a random sample of Sardinian inhabitants. *ATS-SARDEGNA Research Group. Eur J Epidemiol* 12(1):71–76
- Podolin DA, Wills BK, Wood IO, Lopez M, Mazzeo RS, Roth DA (2001) Attenuation of age-related declines in glucagon-mediated signal transduction in rat liver by exercise training. *Am J Physiol* 281(3):E516–E523
- Praphatsorn P, Thong-Ngam D, Kulaputana O, Klaiakaw N (2010) Effects of intense exercise on biochemical and histological changes in rat liver and pancreas. *Asian Biomed* 4(4):619–625
- Prior SJ, Jenkins NT, Brandauer J, Weiss EP, Hagberg JM (2012) Aerobic exercise training increases circulating insulin-like growth factor binding protein-1 concentration, but does not attenuate the reduction in circulating insulin-like growth factor binding protein-1 after a high-fat meal. *Metabolism* 61(3):310–316
- Promrat K, Kleiner DE, Niemeier HM, Jackvony E, Kearns M, Wands JR, Fava J, Wing RR (2010) Randomized controlled trial testing the effect of weight loss on nonalcoholic steatohepatitis (NASSH). *Hepatology* 51(1):121–129
- Radak Z, Chung HY, Naito H, Takahashi K, Jung KJ, Kim HJ, Goto S (2004) Age-associated increase in oxidative stress and nuclear factor B activation are attenuated in rat liver by regular exercise. *FASEB J* 18(6):749–750
- Rama R, Ibáñez J, Riera M, Prats MT, Pagés T, Palacios L (1994) Hematological, electrolyte, and biochemical alterations after a 100-km run. *Can J Appl Physiol* 19(4):411–420
- Rao R (2009) Endotoxemia and gut barrier disease in alcoholic liver disease. *Hepatology* 50:638–644
- Rector SR, Thyfault JP (2011) Does physical inactivity cause nonalcoholic fatty liver disease? *J Appl Physiol* 111(6):1828–1835
- Rector RS, Thyfault JP, Laye MJ, Morris RT, Borengasser SJ, Uptergrove GM, Chakravarthy MV, Booth FW, Ibdah JA (2008) Cessation of daily exercise dramatically alters precursors of hepatic steatosis in Otsuka Long-Evans Tokushima Fatty (OLETF) rats. *J Physiol* 586(Pt 17):4241–4249
- Rector RS, Uptergrove GM, Morris EM, Borengasser SJ, Laughlin MH, Booth FW, Thyfault JP, Ibdah JA (2011) Daily exercise vs. caloric restriction for prevention of nonalcoholic fatty liver disease in the OLETF rat model. *Am J Physiol* 300(5):G874–G883
- Reddy Avula CP, Ferandes G (1999) Modulation of antioxidant enzymes and lipid peroxidation in salivary gland and other tissues in mice by moderate treadmill exercise. *Aging (Milano)* 11(4):246–252
- Reynoso E, Lavine JE (2012) NAFLD: the role of exercise in treating NAFLD. *Nat Rev Gastroenterol Hepatol* 9(7):368–370
- Richards D, Richards R, Schofield PJ, Ross V (1979) Biochemical and haematological changes in Sydney's the sun city-to-surf fun runners. *Med J Austral* 2(9):449–453
- Ringseis R, Mooren FC, Keller J, Couturier A, Wen G, Hirche F, Stangl GI, Eder K, Kruger K (2011) Regular endurance exercise improves the diminished hepatic carnitine status in mice fed a high-fat diet. *Mol Nutr Food Res* 55(Suppl 2):S193–S202
- Ritland S, Foss NE, Gjone E (1982) Physical activity and liver disease and liver function. *Scand J Soc Med* 29:221–226
- Ritland S, Petlund CF, Knudsen T, Skrede S (1983) Improvement of physical capacity after long-term training in patients with chronic active hepatitis. *Scand J Gastroenterol* 18(8):1083–1087
- Robinson D, Whitehead TP (1989) Effect of body mass and other factors on serum liver enzyme levels in men attending for well population screening. *Ann Clin Biochem* 26(Pt 5):393–400
- Rodriguez B, Torres DM, Harrison SA (2012) Physical activity: an essential component of lifestyle modification in NAFLD. *Nat Rev Gastroenterol Hepatol* 9(12):726–731
- Roecker L, Meier-Buttermilch R, Brechtel L, Nemeth E, Ganz T (2005) Iron-regulatory protein hepcidin is increased in female athletes after a marathon. *Eur J Appl Physiol* 95(5–6):569–571
- Rohrer TR, Rizzo VF, Casar JJ, Muelbrecht O, Sprengart S, Gortner L, Stierkorb E (2008) Changes in hepatic risk factors, metabolic variables, body composition, and physical fitness in obese children after a one-year weight loss program. *J Pediatr Endocrinol* 21(9):837–845
- Romani R, De Medio GE, di Tullio S, Lapalombella R, Pirisinu I, Margonato V, Veicsteinas A, Marini M, Rosi G (2009) Modulation of paraoxonase 1 and 3 expression after moderate exercise training in the rat. *J Lipid Res* 50(10):2036–2045
- Romijn JA, Coyle EF, Sidossis LS, Gastaldelli A, Horowitz JF, Endert E, Wolfe RR (1993) Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am J Physiol* 265(3 Pt 1):E380–E391
- Ropelle ER, Pauli JR, Cintra DE, Frederico MJ, de Pinho RA, Velloso LA, De Souza CT (2009) Acute exercise modulates the Foxo1/PGC-1alpha pathway in the liver of diet-induced obesity rats. *J Physiol* 587(Pt 9):2069–2076
- Rosales XQ, Chu ML, Shilling C, Wall C, Pastores GM, Mendell JR (2008) Fidelity of gamma-glutamyl transferase (GGT) in differentiating skeletal muscle from liver damage. *J Child Neurol* 23(7):748–751
- Rowell LB (1971) Visceral blood flow and metabolism during exercise. In: Shephard RJ (ed) *Frontiers of fitness*. CC Thomas, Springfield, pp 210–232
- Rowell LB (1974) Human cardiovascular adjustments to exercise and thermal stress. *Physiol Rev* 54(1):75–159
- Rowell LB (1986) *Human circulation: regulation during physical stress*. Oxford University, New York
- Rowell LB, Blackmon JR, Bruce RA (1964) Indocyanine green clearance and estimated hepatic blood flow during mild to maximal exercise in upright man. *J Clin Invest* 43:1677–1690
- Saengsirisuwan V, Phadungkij S, Pholpramool C (1998) Renal and liver functions and muscle injuries in Thai boxers during training and after competition. *Br J Sports Med* 32(4):304–308
- Salo DC, Donovan CM, Davies KJ (1991) HSP70 and other possible heat shock or oxidative stress proteins are induced in muscle, heart and liver during exercise. *Free Radic Biol Med* 11(3):239–246
- Santomauro M, Paoli-Valeri M, Fernandez M, Camacho N, Molina Z, Cicchetti R, Valeri L, Davila de Campagnaro E, Arata-Bella-barba G (2012) Enfermedad del hígado graso no alcohólico y su asociación con variables clínicas y bioquímicas en los niños y adolescentes obesos: efecto de una intervención de un año en el estilo de vida. [Non-alcoholic fatty liver disease and its association with clinical and biochemical variables in obese children and adolescents: effect of a one-year intervention on lifestyle]. *Endocrinol Nutr* 59(6):346–353
- Schäfer S, Kantartzis K, Machann J, Venter C, Niess A, Schick F, Machicao F, Häring HU, Fritsche A, Stefan N (2007) Lifestyle

- intervention in individuals with normal versus impaired glucose tolerance. *Eur J Clin Invest* 37(7):535–543
- Schoemaker RC, Burggraaf J, Cohen AF (1998) Assessment of hepatic blood flow using continuous infusion of high clearance drugs. *Br J Clin Pharmacol* 45(5):463–469
- Schultz A, Mendonca LS, Aguila MB, Mandarim-de-Lacerda CA (2012) Swimming training beneficial effects in a mice model of nonalcoholic fatty liver disease. *Exp Toxicol Pathol* 64(4):273–282
- Schwarz AJ, Brasel JA, Hintz RL, Mohan S, Cooper DM (1996) Acute effect of brief low- and high-intensity exercise on circulating insulin-like growth factor (IGF) I, II, and IGF-binding protein-3 and its proteolysis in young healthy men. *J Clin Endocrinol Metab* 81(10):3492–3497
- Sen CK, Marin E, Kretzschmar M, Hanninen O (1992) Skeletal muscle and liver glutathione homeostasis in response to training, exercise, and immobilization. *J Appl Physiol* 73(4):1265–1272
- Sene-Fiorese M, Duarte FO, Scarmagnani FRR, Cheik NC, Manzoni MSJ, Nonaka KO, Rossi EA, De Oliveira Duarte ACG, Damaso AR (2009) Efficiency of intermittent exercise on adiposity and fatty liver in rats fed with high-fat diet. *Obesity (Silver Spring)* 16(10):2217–2222
- Seppala-Lindroos A, Vehkavaara S, Hakkinen AM, Goto T, West-erbacka J, Sovijari A, Halavaara J, Yki-Jarvinen H (2007) Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. *J Clin Endocrinol Metab* 87(7):3023–3028
- Shah K, Stufflebam A, Hilton TN, Sinacore DR, Klein S, Villareal DT (2009) Diet and exercise interventions reduce intrahepatic fat content and improve insulin sensitivity in obese older adults. *Obesity (Silver Spring)* 17(12):2162–2168
- Shapiro Y, Magazanik A, Sohar E, Reich CB (1973) Serum enzyme changes in untrained subjects following a prolonged march. *Can J Physiol Pharmacol* 51(4):271–276
- Shephard RJ (1982) *Physiology and biochemistry of exercise*. Praeger Publications, New York
- Shephard RJ (2013) Are the abdominal viscera important to the sport physician? Esophagus, stomach, intestines, spleen and pancreas. In: Shephard RJ et al (eds) *Year book of sports medicine*. Elsevier, Philadelphia
- Shojaee-Moradie F, Baynes KC, Pentecost C, Bell JD, Thomas EL, Jackson NC, Stolinski M, Whyte M, Lovell D, Bowes SB, Gibney J, Jones RH, Umpleby AM (2007) Exercise training reduces fatty acid availability and improves the insulin sensitivity of glucose metabolism. *Diabetologia* 50(2):404–413
- Slentz CA, Bateman LA, Willis LH, Shields AT, Tanner CJ, Piner LW, Hawk VH, Muehlbauer MJ, Samsa GP, Nelson RC, Huffman KM, Bales CW, Houmard JA, Kraus WE (2012) Effects of aerobic vs. resistance training on visceral and liver fat stores, liver enzymes, and insulin resistance by HOMA in overweight adults from STRRIDE AT/RT. *Am J Physiol* 301(5):E1033–E1039
- Smith JE, Garbutt G, Lopes P, Tunstall Pedoe D (2004) Effects of prolonged strenuous exercise (marathon running) on biochemical and haematological markers used in the investigation of patients in the emergency department. *Br J Sports Med* 38(3):292–294
- Socha P, Horvath A, Vajro P, Dziechciarz P, Dhawan A, Szajewska H (2009) Pharmacological interventions for non-alcoholic fatty liver disease in adults and in children: a systematic review. *J Pediatr Gastroenterol Nutr* 48(5):587–596
- Spassiani NA, Kuk JL (2008) Exercise and the fatty liver. *Appl Physiol Nutr Metab* 33(4):802–807
- St George A, Bauman A, Johnston A, Farrell G, Chey T, George J (2009) Independent effects of physical activity in patients with nonalcoholic fatty liver disease. *Hepatology* 50(1):68–76
- Stanca CM, Bach N, Krause C, Tandon N, Freni MA, Gutierrez JA, Bodian C, Lopez J, Berk PD, Bodenheimer HC, Branch AD, Odin JA (2005) Evaluation of fatigue in US patients with primary biliary cirrhosis. *Am J Gastroenterol* 100(5):1104–1109
- Steenivasa Baba C, Alexander G, Kalyani B, Pandey R, Rastogi S, Pandey A, Choudhuri G (2006) Effect of exercise and dietary modification on serum aminotransferase levels in patients with nonalcoholic steatohepatitis. *Gastroenterol Hepatol* 21(1 Pt 1):191–198
- Straznicki NE, Lambert EA, Grima MT, Eikelis N, Nestel PJ, Dawood T, Schlaich MP, Masuo K, Chopra R, Sari CI, Dixon JB, Tilbrook AJ, Lambert GW (2012) The effects of dietary weight loss with or without exercise training on liver enzymes in obese metabolic syndrome subjects. *Diabetes Obes Metab* 14(2):139–148
- Suh SH, Paik IY, Jacobs K (2007) Regulation of blood glucose homeostasis during prolonged exercise. *Mol Cells* 23(3):272–279
- Sullivan S, Kirk E, Mittendorfer B, Patterson B, Klein S (2012) Randomized trial of exercise effect on intrahepatic triglyceride content and lipid kinetics in nonalcoholic fatty liver disease. *Hepatology* 55(6):1738–1745
- Sumida KD, Donovan CM (1995) Enhanced hepatic gluconeogenic capacity for selected precursors after endurance training. *J Appl Physiol* 79(6):1883–1888
- Sumida KD, Urdiales JH, Donovan CM (1993) Enhanced gluconeogenesis from lactate in perfused livers after endurance training. *J Appl Physiol* 74(2):782–787
- Suzuki A, Lindor K, St Saver J, Lymp J, Mendes F, Muto A, Okada T, Angulo P (2005) Effect of changes on body weight and lifestyle in nonalcoholic fatty liver disease. *J Hepatol* 43(6):1060–1066
- Suzuki K, Peake J, Nosaka K, Okutsu M, Abbiss CR, Surriano R, Bishop D, Quod MJ, Lee H, Martin DT, Laursen PB (2006) Changes in markers of muscle damage, inflammation and HSP70 after an Ironman triathlon race. *Eur J Appl Physiol* 98(6):525–534
- Swartz RD, Sidell FR, Cucinell SA (1974) Effects of physical stress on the disposition of drugs eliminated by the liver in man. *J Pharmacol Exp Ther* 188(1):1–7
- Szczepaniak LS, Nurenberg P, Leonard D, Browning JD, Reingold JS, Grundy S, Hobbs HH, Dobbins RL (2005) Magnetic resonance spectroscopy to measure hepatic triglyceride content: prevalence of hepatic steatosis in the general population. *Am J Physiol* 288(2):E462–E468
- Takahashi Y, Fukusato T (2010) Pediatric nonalcoholic fatty liver disease: overview with emphasis on histology. *World J Gastroenterol* 16(42):5280–5285
- Takahashi I, Umeda T, Mashiko T, Chinda D, Oyama T, Sugawara K, Nkaii S (2007) Effects of rugby sevens matches on human neutrophil-related nonspecific immunity. *Br J Sports Med* 41(1):13–18
- Takeshita H, Horiuchi M, Izumo K, Kawaguchi H, Arimura E, Aoyama K, Takeuchi T (2012) Long-term voluntary exercise, representing habitual exercise, lowers visceral fat and alters plasma amino acid levels in mice. *Env Health Prev Med* 17(4):275–284
- Tamura Y, Tanaka Y, Sato F, Choi JB, Watada H, Niwa M, Kinoshita J, Ooka A, Kumashiro N, Igarashi Y, Kyogoku S, Maehara T, Kawasumi M, Hirose T, Kawamori R (2005) Effects of diet and exercise on muscle and liver intracellular lipid contents and insulin sensitivity in type 2 diabetic patients. *J Clin Endocrinol Metab* 90(6):3191–3196
- Targher GB, Bellis A, Fornengo P, Ciaravella F, Pichiri I, Cavallo Perin P, Trimarco B, Marchesini G (2010) Prevention and treatment of nonalcoholic fatty liver disease. *Dig Liver Dis* 42(5):331–340
- Tarter RE, Panzak G, Switala J, Lu S, Simkevitz HV, Thiel D (1997) Isokinetic muscle strength and its association with

- neuropsychological capacity in cirrhotic alcoholics. *Alcohol Clin Exp Res* 21(2):191–196
- Terao T, Fujise T, Uchiyama S, Yamashita Y, Nakano S (1989) Effects of swimming exercise at two different water temperature on hepatic lipid and lipoprotein levels in experimental fatty liver rats. *Tokai J Exp Clin Med* 14(2):139–145
- Terjung RL, Baldwin KM, Winder WW, Holloszy JO (1971) Glycogen repletion in different types of muscle and in liver after exhausting exercise. *Am J Physiol* 226(6):1387–1391
- Terziyski K, Andonov V, Marinov B, Kostianev S (2008) Exercise performance and ventilatory efficiency in patients with mild and moderate liver cirrhosis. *Clin Exp Pharmacol Physiol* 35(2):135–140
- Thamer C, Machann J, Stefan N, Haap M, Schäfer S, Brenner S, Kantartzis K, Claussen C, Schick F, Haring H, Fritsche A (2007) High visceral fat mass and high liver fat are associated with resistance to lifestyle intervention. *Obesity (Silver Spring)* 15(2):531–538
- Thamer C, Jürgen M, Stefan N, Schäfer SA, Machicao F, Staiger H, Laakso M, Böttcher M, Claussen C, Schick F, Fritsche A, Haring H-U (2008) Variations in PPARD determine the change in body composition during lifestyle intervention: a whole body magnetic resonance study. *J Clin Endocrinol Metab* 93(4):1497–1500
- Theilade P, Hansen JM, Skovsted L, Kampmann JP (1979) Effect of exercise on thyroid parameters and on metabolic clearance rate of antipyrine in man. *Acta Endocrinol* 92(2):271–276
- Thoma C, Day CP, Trenell MI (2012) Lifestyle interventions for the treatment of non-alcoholic fatty liver disease in adults: a systematic review. *J Hepatol* 56(1):255–266
- Thomas EL, Brynes AE, Hamilton G, Patel N, Spong A, Goldin RD, Frost G, Bell JD, Taylor-Robinson SD (2006) Effect of nutritional counselling on hepatic, muscle and adipose tissue fat content and distribution in non-alcoholic fatty liver disease. *World J Gastroenterol* 12(36):5813–5819
- Thompson WR (2010) ACSM's guidelines for exercise testing and prescription, 8th ed, Lippincott, Williams & Wilkins, Baltimore
- Thompson PD, Cullinane EM, Sady SP, Flynn MM, Chenevert CB, Herbert PN (1991) High-density-lipoprotein metabolism in endurance athletes and sedentary men. *Circulation* 84(1):140–152
- Thompson PD, Crouse SF, Goodpaster B, Kelley D, Moyna N, Pescatello L (2001) The acute versus the chronic response to exercise. *Med Sci Sports Exerc* 33(6 Suppl):S438–S445 (discussion S452–S453)
- Thompson D, Markovitch D, Betts JA, Mazzatti D, Turner J, Tyrrell RM (2010) Time course of changes in inflammatory markers during a 6-mo exercise intervention in sedentary middle-aged men: a randomized controlled trial. *J Appl Physiol* 108(4):769–779
- Thyfault JP, Rector RS, Uptergrove GM, Borengasser SJ, Morris EM, Wei Y, Laye MJ, Burant CF, Qi NR, Ridenour SE, Koch LG, Britton SL, Ibdah JA (2009) Rats selectively bred for low aerobic capacity have reduced hepatic mitochondrial capacity and susceptibility to hepatic steatosis and injury. *J Physiol* 587(Pt 8):1805–1816
- Tiikkainen M, Beerholm R, Hakkinen AM, Bergholm R, Vehkavaara S, Halavaara J, Teramo K, Rissanen A, Yki-Järvinen H (2002) Liver fat accumulation and insulin resistance in obese women with previous gestational diabetes. *Obesity Res* 10(9):859–867
- Trimmer JK, Schwarz JM, Cadsazza GA, Horning MA, Rodriguez N, Brooks GA (2002) Measurements of gluconeogenesis in exercising men by mass isotopomer distribution analysis. *J Appl Physiol* 93(1):233–241
- Tsekouras YE, Magkos F, Kellas Y, Basioukas KN, Kavouras SA, Sidossis LS (2008) High-intensity interval aerobic training reduces hepatic very low-density lipoprotein-triglyceride secretion rate in men. *Am J Physiol* 295(4):E851–E858
- Tsuruta G, Tanaka N, Hongo M, Komatsu M, Horiuchi A, Hamamoto K, Iguchi C, Nakayama Y, Umemura T, Ichijo T, Matsumoto A, Yoshizawa K, Aoyama T, Tanaka E (2010) Nonalcoholic fatty liver disease in Japanese junior high school students: its prevalence and relationship to lifestyle habits. *J Gastroenterol* 45(6):666–672
- Turgut G, Demir S, Genc O, Karabulut I, Akalin N (2003) The effect of swimming exercise on lipid peroxidation in the rat brain, liver and heart. *Acta Physiol Pharmacol Bulg* 27(2–3):43–45
- Turner JE, Hodges NJ, Bosch JA, Aldred S (2011) Prolonged depletion of antioxidant capacity after ultraendurance exercise. *Med Sci Sports Exerc* 43(9):1770–1776
- Ueno T, Sugawara H, Sujaku K, Hashimoto O, Tsuji R, Tamaki S, Torimura T, Inuzuka S, Sata M, Tanikawa K (1997) Therapeutic effect of restricted diet and exercise in obese patients with fatty liver. *J Hepatol* 27(1 Pt 1):103–107
- van der Heijden G, Toffolo G, Manesso E, Sauer PJ, Sunehag AL (2009) Aerobic exercise increases peripheral and hepatic insulin sensitivity in sedentary adolescents. *J Clin Endocrinol Metab* 94(11):4292–4299
- van der Heijden GJ, Wang ZJ, Chu Z, Toffolo G, Manesso E, Sauer PJ, Sunehag AL (2010a) Strength exercise improves muscle mass and hepatic insulin sensitivity in obese youth. *Med Sci Sports Exerc* 42(11):1973–1980
- van der Heijden GJ, Wang ZJ, Chu ZD, Sauer PJ, Haymond MW, Rodriguez LM, Sunehag AL (2010b) A 12-week aerobic exercise program reduces hepatic fat accumulation and insulin resistance in obese, Hispanic adolescents. *Obesity (Silver Spring)* 18(2):384–390
- van Dijk B, Balkan B, Lindfeldt J, Bouws G, Scheurink AJ, Ahren B, Steffens AB (1994) Contribution of liver nerves, glucagon and adrenaline to the glycemic response to exercise in rats. *Acta Physiol Scand* 150(3):305–313
- van Griensven JM, Burggraaf KJ, Gerloff J, Gunzler WA, Beier H, Kroon R, Huisman LG, Schoemaker RC, Klufft K, Cohen AF (1995) Effects of changing liver blood flow by exercise and food on kinetics and dynamics of saruplase. *Clin Pharmacol Ther* 57(4):381–389
- Van Rensburg JP, Kielblock AJ, van der Linde A (1986) Physiologic and biochemical changes during a triathlon competition. *Int J Sports Med* 7(1):30–35
- van Wijck K, Lenaerts K, van Loon LJC, Peters WHM, Buurman WS, Dejong CHC (2011) Exercise-induced splanchnic hypoperfusion results in gut dysfunction in healthy men. *PLoS One* 6(7):e22366
- Venditti P, Di Meo S (1997) Effect of training on antioxidant capacity, tissue damage, and endurance of adult male rats. *Int J Sports Med* 18(7):497–502
- Viitasalo A, Laaksonen DE, Lindi V, Eloranta AM, Jaaskelainen J, Tompuri T, Vaisanen S, Lakka HM, Lakka TA (2012) Clustering of metabolic risk factors is associated with high-normal levels of liver enzymes among 6- to 8-year-old children: the PANIC study. *Metab Syndr Relat Disord* 10(5):337–343
- Villa JG, Cuadrado G, Bayón JE, González-Gallego J (1998) The effect of physical conditioning on antipyrine clearance. *Eur J Appl Physiol* 77(1–2):106–111
- Wade OL, Bishop JM (1962) Cardiac output and regional blood flow. Blackwell Scientific, Oxford
- Wahren DG, Ekberg K (2007) Splanchnic regulation of glucose production. *Annu Rev Nutr* 27:329–345
- Wahren J, Felig P, Ahlborg G, Jorfeldt L (1971) Glucose metabolism during leg exercise in man. *J Clin Invest* 50(12):2715–2725
- Waskiewicz Z, Klapińska B, Sadowska-Krepa E, Czuba M, Kempa K, Kimsa E, Gerasimuk D, Waskiewicz Z, Klapińska B,

- Sadowska-Krepa E, Czuba M, Kempa K, Kimsa E, Gerasimuk D (2012) Acute metabolic responses to a 24-h ultra-marathon race in male amateur runners. *Eur J Appl Physiol* 112(5):1679–1688
- Wasserman DH, Cherrington AD (1991) Hepatic fuel metabolism during muscular work: role and regulation. *Am J Physiol* 260(6 Pt 1):E811–E824
- Wasserman DH, Spalding JA, Lacy DB, Colburn CA, Goldstein RE, Cherrington AD (1989) Glucagon is a primary controller of hepatic glycogenolysis and gluconeogenesis during muscular work. *Am J Physiol* 257(1 Pt 1):E108–E117
- Wasserman DH, Connolly CC, Pagliassotti MJ (1991) Regulation of hepatic lactate balance during exercise. *Med Sci Sports Exerc* 23(8):912–919
- Wasserman DH, Lacy DB, Bracy DP (1993) Relationship between arterial and portal vein immunoreactive glucagon during exercise. *J Appl Physiol* 75(2):724–729
- Wasserman DH, O'Doherty RM, Zinker BA (1995) Role of the endocrine pancreas in control of fuel metabolism by the liver during exercise. *Int J Obesity Relat Metab Disord* 19(Suppl 4):S22–S30
- Whitfield JB (2001) Gamma glutamyl transferase. *Crit Rev Clin Lab Sci* 38(4):263–355
- Wieczorek-Baranowska A, Nowak A, Michalak E, Karolkiewicz J, Pospieszna B, Rutkowski R, Laurentowska M, Pilaczyńska-Szcześniak L (2011) Effect of aerobic exercise on insulin, insulin-like growth factor-1 and insulin-like growth factor binding protein-3 in overweight and obese postmenopausal women. *J Sports Med Phys Fitness* 51(3):525–532
- Wiesinger GF, Quittan M, Zimmermann K, Nuhr M, Wichlas M, Bodingerbauer M, Asari R, Berlakovich G, Crevenna R, Fialka-Moser V, Peck-Radosavljevic M (2001) Physical performance and health-related quality of life in men on a liver transplantation waiting list. *J Rehabil Med* 33(6):260–265
- Williams P, Wood P, Haskell W, Vranizan K, Ho P, Vodak P (1981) Effect of exercise intensity and duration on plasma-lipoprotein cholesterol levels. *Circulation* 64(4):185
- Wilson PO, Johnson P (2000) Exercise modulates antioxidant enzyme gene expression in rat myocardium and liver. *J Appl Physiol* 88(5):1791–1796
- Winnick JJ, Sherman WM, Habash DL, Stout MB, Failla ML, Belury MA, Schuster DP (2008) Short-term aerobic exercise training in obese humans with type 2 diabetes mellitus improves whole-body insulin sensitivity through gains in peripheral, not hepatic insulin sensitivity. *J Clin Endocrinol Metab* 93(3):771–778
- Wolfe BM, Havel JR, Marliss EB, Kane JP, Seymour J, Ahuja SP (1976) Effects of a 3-day fast and of ethanol on splanchnic metabolism of ffa, amino-acids, and carbohydrates in healthy young men. *J Clin Invest* 57(2):329–340
- Wolfe RR, Klein S, Carraro F, Weber JM (1990) Role of triglyceride fatty-acid cycle in controlling fat-metabolism in humans during and after exercise. *Am J Physiol* 258(2):E382–E389
- Wong F, Girgrah N, Graba J, Allidina Y, Liu P, Blendis L (2001) The cardiac response to exercise in cirrhosis. *Gut* 49(2):268–275
- Woodruff TK, Krummen L, Chen SA, Lyon R, Hansen SE, DeGuzman G, Covello R, Mather J, Cossum P (1993) Pharmacokinetic profile of recombinant human (rh) inhibin A and activin A in the immature rat. II. Tissue distribution of [125I]rh-inhibin A and [125I]rh-activin A in immature female and male rats. *Endocrinology* 132(2):725–734
- Wu HJ, Chen KT, Shee BW, Chang HC, Huang YJ, Yang RS (2004) Effects of 24 h ultra-marathon on biochemical and hematological parameters. *World J Gastroenterol* 10(18):2711–2714
- Wu LJ, Wu MS, Lien GS, Chen FC, Tsai JC (2012) Fatigue and physical activity levels in patients with liver cirrhosis. *J Clin Nurs* 21(1–2):129–138
- Yang RC, Mack GW, Wolfe RR, Nadel ER (1998) Albumin synthesis after intense intermittent exercise in human subjects. *J Appl Physiol* 84(2):584–592
- Yano L, Yano H, Taketa K (1996) Electromagnetic determination of portal venous flow in rats during exercise. *Int Hepatol Comm* 5(3):184–190
- Yano H, Kinoshita S, Yano L (1997) Acute exercise induces mitochondria swelling of hepatocytes surrounding the terminal hepatic venule in rat liver acinus. *Jpn J Phys Fitness Sports Med* 46(1):49–54
- Yasari S, Paquette A, Charbonneau A, Gauthier MS, Savard R, Lavoie JM (2006) Effects of ingesting a high-fat diet upon exercise-training cessation on fat accretion in the liver and adipose tissue of rats. *Appl Physiol Nutr Metab* 31(4):367–375
- Yasari S, Wang D, Prud'homme D, Jankowski M, Gutkowska J, Lavoie JM (2009) Exercise training decreases plasma leptin levels and the expression of hepatic leptin receptor-a, -b, and, -e in rats. *Mol Cell Biochem* 324(1–2):13–20
- Yasari S, Prud'homme D, Wang D, Jankowski M, Levy E, Gutkowska J, Lavoie J-M (2010) Exercise training decreases hepatic SCD-1 gene expression and protein content in rats. *Mol Cell Biochem* 335(1–2):291–299
- Yiamouyiannis CA, Sanders RA, Watkins JB, Martin BJ (1992) Chronic physical activity: hepatic hypertrophy and increased total biotransformation enzyme activity. *Biochem Pharmacol* 44(1):121–127
- Yiamouyiannis CA, Martin BJ, Walkins JB (1993) Chronic physical activity alters excretory function in rats. *J Pharmacol Exp Ther* 265(1):321–327
- Yoneda MF, Fujita K, Iwasaki T, Maeyama S, Terauchi Y, Nakajima A (2006) Treatment of NASH: nutritional counseling and physical exercise. *Nihon Rinsho* 64(6):1139–1145
- Zawalich W, Matuso S, Felig P (1982) Influence of physical training on insulin release and glucose utilization by islet cells. *Am J Physiol* 243(6):E464–E469
- Zelber-Sagi S, Nitzan-Kaluski D, Goldsmith R, Webb M, Zvibel I, Goldiner I, Blendis L, Halpern Z, Oren R (2008) Role of leisure-time physical activity in nonalcoholic fatty liver disease: a population-based study. *Hepatology* 48(6):1791–1798
- Zelber-Sagi S, Ratzu V, Oren R (2011) Nutrition and physical activity in NAFLD: an overview of the epidemiological evidence. *World J Gastroenterol* 17(29):3377–3389
- Zucker DM (2004) An exercise intervention to prevent hepatitis related fatigue. *Hepatology* 40:511A

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.